



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

|   |  |   |
|---|--|---|
| <p>(51) International Patent Classification <sup>6</sup> :<br/>C09B 23/14, C07C 65/19, 65/28, 69/88,<br/>C07F 7/22, C07B 59/00, G01N 33/68,<br/>A61K 31/60, 49/00, 51/00 // C07D<br/>209/48, 261/20, 237/32, 265/02, 265/26,<br/>217/24, 221/14, 257/04, 271/10</p>   | A1   | <p>(11) International Publication Number: <b>WO 98/47969</b></p> <p>(43) International Publication Date: 29 October 1998 (29.10.98)</p> |
| <p>(21) International Application Number: PCT/US98/07889</p> <p>(22) International Filing Date: 20 April 1998 (20.04.98)</p> <p>(30) Priority Data:<br/>08/837,494 18 April 1997 (18.04.97) US</p> <p>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application<br/>US 08/837,494 (CIP)<br/>Filed on 18 April 1997 (18.04.97)</p> <p>(71) Applicant (for all designated States except US): UNIVERSITY OF PITTSBURGH [US/US]; 911 William Pitt Union, Pittsburgh, PA 15260 (US).</p> <p>(72) Inventors; and<br/>(75) Inventors/Applicants (for US only): KLUNK, William, E. [US/US]; 1108 Onondago Street, Pittsburgh, PA 15218 (US). PETTEGREW, Jay, W. [US/US]; 630 S. Linden Avenue, Pittsburgh, PA 15208 (US). MATHIS, Chester, A.,</p>  | <p>Jr. [US/US]; 48 Fox Pionte Drive, Pittsburgh, PA 15238 (US).</p> <p>(74) Agents: BENT, Stephen, A. et al.; Foley &amp; Lardner, Suite 500, 3000 K Street, N.W., Washington, DC 20007-5109 (US).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p><b>Published</b><br/><i>With international search report.<br/>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p> |   |
| <p>(54) Title: ALKYL, ALKENYL AND ALKYNYL CHRYSAMINE G DERIVATIVES FOR THE ANTEMORTEM DIAGNOSIS OF ALZHEIMER'S DISEASE AND <i>IN VIVO</i> IMAGING AND PREVENTION OF AMYLOID DEPOSITION</p> <p>(57) Abstract</p> <p>Amyloid binding compounds which are non-azo derivatives of Chrysamine G, pharmaceutical compositions containing, and methods using such compounds to identify Alzheimer's brain <i>in vivo</i> and to diagnose other pathological conditions characterized by amyloidosis, such as Down's Syndrome are described. Pharmaceutical compositions containing non-azo derivatives of Chrysamine G and methods using such compositions to prevent cell degeneration and amyloid-induced toxicity in amyloidosis associated conditions are also described. Methods using non-azo Chrysamine G derivatives to stain or detect amyloid deposits in biopsy or post-mortem tissue are also described. Methods using non-azo Chrysamine G derivatives to quantify amyloid deposits in homogenates of biopsy and post-mortem tissue are also described.</p> |  |   |

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

|    |                          |    |  |    |  |    |                          |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania                  | ES | Spain                                    | LS | Lesotho                                      | SI | Slovenia                 |
| AM | Armenia                  | FI | Finland                                  | LT | Lithuania                                    | SK | Slovakia                 |
| AT | Austria                  | FR | France                                   | LU | Luxembourg                                   | SN | Senegal                  |
| AU | Australia                | GA | Gabon                                    | LV | Latvia                                       | SZ | Swaziland                |
| AZ | Azerbaijan               | GB | United Kingdom                           | MC | Monaco                                       | TD | Chad                     |
| BA | Bosnia and Herzegovina   | GE | Georgia                                  | MD | Republic of Moldova                          | TG | Togo                     |
| BB | Barbados                 | GH | Ghana                                    | MG | Madagascar                                   | TJ | Tajikistan               |
| BE | Belgium                  | GN | Guinea                                   | MK | The former Yugoslav<br>Republic of Macedonia | TM | Turkmenistan             |
| BF | Burkina Faso             | GR | Greece                                   | ML | Mali   | TR | Turkey                   |
| BG | Bulgaria                 | HU | Hungary                                  | MN | Mongolia                                     | TT | Trinidad and Tobago      |
| BJ | Benin                    | IE | Ireland                                  | MR | Mauritania                                   | UA | Ukraine                  |
| BR | Brazil                   | IL | Israel                                   | MW | Malawi                                       | UG | Uganda                   |
| BY | Belarus                  | IS | Iceland                                  | MX | Mexico                                       | US | United States of America |
| CA | Canada                   | IT | Italy                                    | NE | Niger  | UZ | Uzbekistan               |
| CF | Central African Republic | JP | Japan                                    | NL | Netherlands                                  | VN | Viet Nam                 |
| CG | Congo                    | KE | Kenya                                    | NO | Norway                                       | YU | Yugoslavia               |
| CH | Switzerland              | KG | Kyrgyzstan                               | NZ | New Zealand                                  | ZW | Zimbabwe                 |
| CI | Côte d'Ivoire            | KP | Democratic People's<br>Republic of Korea | PL | Poland                                       |    |                          |
| CM | Cameroon                 | KR | Republic of Korea                        | PT | Portugal                                     |    |                          |
| CN | China                    | KZ | Kazakhstan                               | RO | Romania                                      |    |                          |
| CU | Cuba                     | LC | Saint Lucia                              | RU | Russian Federation                           |    |                          |
| CZ | Czech Republic           | LI | Liechtenstein                            | SD | Sudan  |    |                          |
| DE | Germany                  | LK | Sri Lanka                                | SE | Sweden                                       |    |                          |
| DK | Denmark                  | LR | Liberia                                  | SG | Singapore                                    |    |                          |
| EE | Estonia                  |    |  |    |  |    |                          |

**ALKYL, ALKENYL AND ALKYNYL CHRYSAMINE G  
DERIVATIVES FOR THE ANTEMORTEM DIAGNOSIS OF  
ALZHEIMER'S DISEASE AND IN VIVO IMAGING AND  
PREVENTION OF AMYLOID DEPOSITION**

The present invention was made utilizing funds from the National Institute of Ageing, grant numbers AG-05443, AG-05133 and AG-08974. This is a continuation-in-part application of U.S. patent application Serial No. 08/640,740 and PCT/US96/05918, both filed May 1, 1996 which is a continuation-in-part application of U.S. patent application Serial No. 08/432,019, filed May 1, 1995 which is a continuation-in-part of U.S. patent application Serial No. 08/282,289, filed July 29, 1994.

10                   Background of the Invention

The present invention relates to the identification of compounds that are suitable for imaging amyloid deposits in living patients. More specifically, the present invention relates to a method of imaging amyloid deposits in brain *in vivo* to allow antemortem diagnosis of Alzheimer's Disease. The present invention also relates to therapeutic uses for such compounds.

Alzheimer's Disease ("AD") is a neurodegenerative illness characterized by memory loss and other cognitive deficits. McKhann et al., *Neurology* 34: 939 (1984). It is the most common cause of dementia in the United States. AD can strike persons as young as 40-50 years of age. Yet, because the presence of the disease is difficult to determine without dangerous brain biopsy, the time of onset is unknown. The prevalence of AD increases with age, with estimates of the affected

-2-

population reaching as high as 40-50% by ages 85-90.  
Evans et al., *JAMA* 262: 2551 (1989); Katzman, *Neurology*  
43: 13 (1993).

By definition, AD is definitively diagnosed through  
5 examination of brain tissue, usually at autopsy.  
Khachaturian, *Arch. Neurol.* 42: 1097 (1985); McKhann et  
al., *Neurology* 34: 939 (1984). Neuropathologically, this  
disease is characterized by the presence of neuritic  
10 plaques (NP), neurofibrillary tangles (NFT), and neuronal  
loss, along with a variety of other findings. Mann,  
*Mech. Ageing Dev.* 31: 213 (1985). Post-mortem slices of  
brain tissue of victims of Alzheimer's disease exhibit  
the presence of amyloid in the form of proteinaceous  
extracellular cores of the neuritic plaques that are  
15 characteristic of AD.

The amyloid cores of these neuritic plaques are  
composed of a protein called the  $\beta$ -amyloid ( $A\beta$ ) that is  
arranged in a predominately beta-pleated sheet  
configuration. Mori et al., *Journal of Biological*  
20 *Chemistry* 267: 17082 (1992); Kirschner et al., *PNAS* 83:  
503 (1986). Neuritic plaques are an early and invariant  
aspect of the disease. Mann et al., *J. Neurol. Sci.* 89:  
169; Mann, *Mech. Ageing Dev.* 31: 213 (1985); Terry et  
al., *J. Neuropathol. Exp. Neurol* 46: 262 (1987).

25 The initial deposition of  $A\beta$  probably occurs long  
before clinical symptoms are noticeable. The currently  
recommended "minimum microscopic criteria" for the  
diagnosis of AD is based on the number of neuritic  
plaques found in brain. Khachaturian, *Arch. Neurol.*,  
30 *supra* (1985). Unfortunately, assessment of neuritic  
plaque counts must be delayed until after death.

Amyloid-containing neuritic plaques are a prominent  
feature of selective areas of the brain in AD as well as  
Downs Syndrome and in persons homozygous for the  
35 apolipoprotein E4 allele who are very likely to develop  
AD. Corder et al., *Science* 261: 921 (1993); Divry, P.,  
*J. Neurol. Psych.* 27: 643-657 (1927); Wisniewski et al.,

-3-

in Zimmerman, H.M. (ed.): PROGRESS IN NEUROPATHOLOGY,  
(Grune and Stratton, N.Y. 1973) pp. 1-26. Brain  
amyloid is readily demonstrated by staining brain  
sections with thioflavin S or Congo red. Puchtler et  
al., *J. Histochem. Cytochem.* 10: 35 (1962). Congo red  
stained amyloid is characterized by a dichroic  
appearance, exhibiting a yellow-green polarization color.  
The dichroic binding is the result of the beta-pleated  
sheet structure of the amyloid proteins. Glenner, G.  
*N. Eng. J. Med.* 302: 1283 (1980). A detailed discussion  
of the biochemistry and histochemistry of amyloid can be  
found in Glenner, *N. Eng. J. Med.*, 302: 1333 (1980).

Thus far, diagnosis of AD has been achieved mostly  
through clinical criteria evaluation, brain biopsies and  
post mortem tissue studies. Research efforts to develop  
methods for diagnosing Alzheimer's disease *in vivo*  
include (1) genetic testing, (2) immunoassay methods and  
(3) imaging techniques.

Evidence that abnormalities in A $\beta$  metabolism are  
necessary and sufficient for the development of AD is  
based on the discovery of point mutations in the A $\beta$   
precursor protein in several rare families with an  
autosomal dominant form of AD. Hardy, *Nature Genetics* 1:  
233 (1992); Hardy et al., *Science* 256: 184 (1992). These  
mutations occur near the N- and C-terminal cleavage  
points necessary for the generation of A $\beta$  from its  
precursor protein. St. George-Hyslop et al., *Science*  
235: 885 (1987); Kang et al., *Nature* 325: 733 (1987);  
Potter WO 92/17152. Genetic analysis of a large number  
of AD families has demonstrated, however, that AD is  
genetically heterogeneous. St. George-Hyslop et al.,  
*Nature* 347: 194 (1990). Linkage to chromosome 21 markers  
is shown in only some families with early-onset AD and in  
no families with late-onset AD. More recently a gene on  
chromosome 14 whose product is predicted to contain  
multiple transmembrane domains and resembles an integral  
membrane protein has been identified by Sherrington et

-4-

al., *Nature* 375: 754-760 (1995). This gene may account for up to 70% of early-onset autosomal dominant AD.

Preliminary data suggests that this chromosome 14 mutation causes an increase in the production of A $\beta$ .

- 5 Scheuner et al., *Soc. Neurosci. Abstr.* 21: 1500 (1995).  
A mutation on a very similar gene has been identified on chromosome 1 in Volga German kindreds with early-onset AD. Levy-Lahad et al., *Science* 269: 973-977 (1995).

- Screening for apolipoprotein E genotype has been  
10 suggested as an aid in the diagnosis of AD. Scott, *Nature* 366: 502 (1993); Roses, *Ann. Neurol.* 38: 6-14 (1995). Difficulties arise with this technology, however, because the apolipoprotein E4 allele is only a risk factor for AD, not a disease marker. It is absent  
15 in many AD patients and present in many non-demented elderly people. Bird, *Ann. Neurol.* 38: 2-4 (1995).

- Immunoassay methods have been developed for detecting the presence of neurochemical markers in AD patients and to detect an AD related amyloid protein in  
20 cerebral spinal fluid. Warner, *Anal. Chem.* 59: 1203A (1987); World Patent No. 92/17152 by Potter; Glenner et al., U.S. Patent No. 4,666,829. These methods for diagnosing AD have not been proven to detect AD in all patients, particularly at early stages of the disease and  
25 are relatively invasive, requiring a spinal tap. Also, attempts have been made to develop monoclonal antibodies as probes for imaging of A $\beta$ . Majocha et al., *J. Nucl. Med.*, 33: 2184 (1992); Majocha et al., WO 89/06242 and Majocha et al., U.S. Patent 5,231,000. The major  
30 disadvantage of antibody probes is the difficulty in getting these large molecules across the blood-brain barrier. Using antibodies for in vivo diagnosis of AD would require marked abnormalities in the blood-brain barrier in order to gain access into the brain. There is  
35 no convincing functional evidence that abnormalities in the blood-brain barrier reliably exist in AD. Kalaria, *Cerebrovascular & Brain Metabolism Reviews* 4: 226 (1992).

-5-

A $\beta$  antibodies are also disadvantageous for use in AD diagnostics because they typically stain deposits of A $\beta$  containing non- $\beta$ -sheet (non-fibrillar) A $\beta$  in addition to the neuritic plaques. Yamaguchi et al., *Acta Neuropathol.*, 77: 314 (1989). These deposits may be a separate type of lesion, not necessarily involved in the dementing process of AD. The latter is suggested by findings of nonfibrillar amyloid deposits in cognitively normal controls and aged dogs. Moran et al., *Medicina Clinica* 98: 19 (1992); Shimada et al., *Journal of Veterinary Medical Science* 54: 137 (1992); Ishihara et al., *Brain Res.* 548: 196 (1991); Giaccone et al., *Neurosci. Lett.* 114: 178 (1990). Even if non-fibrillar amyloid deposits are forerunners of neuritic plaques, the key pathological event in AD may be the process that turns the apparently benign non-fibrillar amyloid deposit into the neuritic plaque with its associated halo of degeneration. Therefore, a probe is needed that is specific for the fibrillar A $\beta$  deposits and NFTs as a more specific marker for AD pathophysiology than antibodies that would also label non-fibrillar amyloid deposits.

Recently, radiolabeled A $\beta$  peptide has been used to label diffuse, compact and neuritic type plaques in sections of AD brain. Maggio et al., WO 93/04194. However, these peptides share all of the disadvantages of antibodies. Specifically, peptides do not normally cross the blood-brain barrier in amounts necessary for imaging.

Congo red may be used for diagnosing amyloidosis in vivo in non-brain parenchymal tissues. However, Congo red is probably not suitable for in vivo diagnosis of A $\beta$  deposits in brain because only 0.03% of an injected dose of iodinated Congo red can enter the brain parenchyma. Tubis et al., *J. Amer. Pharm. Assn.* 49: 422 (1960). Radioiodinated bisdiazobenzidine compounds related to Congo red, such as Benzo Orange R and Direct Blue 4, have been proposed to be useful in vitro and in vivo to detect the presence and location of amyloid deposits in an organ

-6-

of a patient. Quay et al., U.S. Patent Nos. 5,039,511 and 4,933,156. However, like Congo red, all of the compounds proposed by Quay contain strongly acidic sulfonic acid groups which severely limit entry of these compounds into the brain making it extremely difficult to attain an imaging effective quantity or detectable quantity in the brain parenchyma.

The inability to assess amyloid deposition in AD until after death impedes the study of this devastating illness. A method of quantifying amyloid deposition before death is needed both as a diagnostic tool in mild or clinically confusing cases as well as in monitoring the effectiveness of therapies targeted at preventing A $\beta$  deposition. Therefore, it remains of utmost importance to develop a safe and specific method for diagnosing AD before death by imaging amyloid in brain parenchyma *in vivo*. Even though various attempts have been made to diagnose AD *in vivo*, currently, there are no antemortem probes for brain amyloid. No method has utilized a high affinity probe for amyloid that has low toxicity, can cross the blood-brain barrier, and binds more effectively to AD brain than to normal brain in order to identify AD amyloid deposits in brain before a patient's death. Thus, no *in vivo* method for AD diagnosis has been demonstrated to meet these criteria.

Very recent data suggest that amyloid-binding compounds will have therapeutic potential in AD and type 2 diabetes mellitus. As mentioned above, there are two broad categories of plaques in AD brain, diffuse and neuritic (classical). Diffuse plaques do not appear to induce morphological reactions such as the reactive astrocytes, dystrophic neurites, microglia cells, synapse loss, and full complement activation found in neuritic plaques. Joachim et al., *Am. J. Pathol.* 135: 309 (1989); Masliah et al., *loc. cit.* 137: 1293 (1990); Lue and Rogers, *Dementia* 3: 308 (1992). These morphological reactions all signify that neurotoxic and cell



-7-

degenerative processes are occurring in the areas adjacent to the fibrillar A $\beta$  deposits of neuritic plaques. A $\beta$ -induced neurotoxicity and cell degeneration has been reported in a number of cell types *in vitro*.

- 5 Yankner et al., *Science* 250: 279 (1990); Roher et al., *BBRC* 174: 572 (1991); Frautschy et al., *Proc. Natl. Acad. Sci.* 88: 83362 (1991); Shearman et al., *loc. cit.* 91: 1470 (1994). It has been shown that aggregation of the A $\beta$  peptide is necessary for *in vitro* neurotoxicity.
- 10 Yankner, *Neurobiol. Aging* 13: 615 (1992). Differences in the state of aggregation of A $\beta$  in diffuse and neuritic plaques may explain the lack of neurotoxic response surrounding the diffuse plaque. Lorenzo and Yankner, *Proc. Natl. Acad. Sci.*, 91: 12243 (1994). Recently,
- 15 three laboratories have reported results which suggest that Congo red inhibits A $\beta$ -induced neurotoxicity and cell degeneration *in vitro*. Burgevin et al., *NeuroReport* 5: 2429 (1994); Lorenzo and Yankner, *Proc. Natl. Acad. Sci.* 91: 12243 (1994); Pollack et al., *Neuroscience Letters*
- 20 184: 113 (1995); Pollack et al., *Neuroscience Letters* 197: 211 (1995). The mechanism appears to involve both inhibition of fibril formation and prevention of the neurotoxic properties of formed fibrils. Lorenzo and Yankner, *Proc. Natl. Acad. Sci.* 91: 12243 (1994). Congo
- 25 red also has been shown to protect pancreatic islet cells from the toxicity caused by amylin. Lorenzo and Yankner, *Proc. Natl. Acad. Sci.* 91: 12243 (1994). Amylin is a fibrillar peptide similar to A $\beta$  which accumulates in the pancreas in type 2 diabetes mellitus.

- 30 It is known in the art that certain azo dyes may be carcinogenic. Morgan et al. *Environmental Health Perspectives*, 102(supp.) 2: 63-78, (1994). This potential carcinogenicity appears to be based largely on the fact that azo dyes are extensively metabolized to the
- 35 free parent amine by intestinal bacteria. Cerniglia et al., *Biochem. Biophys. Res. Com.*, 107: 1224-1229, (1982). In the case of benzidine dyes (and many other substituted

-8-

benzidines), it is the free amine which is the carcinogen. These facts have little implication for amyloid imaging studies in which an extremely minute amount of the high specific activity radiolabelled dye would be directly injected into the blood stream. In this case, the amount administered would be negligible and the dye would by-pass the intestinal bacteria.

5 In the case of therapeutic usage, these facts have critical importance. Release of a known carcinogen from a therapeutic compound is unacceptable. A second problem with diazo dye metabolism is that much of the administered drug is metabolized by intestinal bacteria prior to absorption. This lowered bioavailability remains a disadvantage even if the metabolites released are innocuous.

10 Thus, a need exists for amyloid binding compounds which are similar to Congo red but which enter the brain (Congo Red does not). Such compounds could be used in preventing cell degeneration associated with fibril formation and thereby treat pathological conditions in amyloid associated diseases, such as AD and Downs Syndrome and in treating pancreatic islet cell toxicity in type 2 diabetes mellitus.

20 A further need exists for amyloid binding compounds that are non-toxic and bioavailable and, consequently, can be used in therapeutics.

#### Summary of the Invention

It therefore is an object of the present invention to provide a safe, specific method for diagnosing AD before death by in vivo imaging of amyloid in brain parenchyma. It is another object of the present invention to provide an approach for identifying AD amyloid deposits in brain before a patient's death, using a high-affinity probe for amyloid which has low toxicity, can cross the blood-brain barrier, and can distinguish AD brain from normal brain. It is another object to provide

(I)

The figure displays eight chemical structures under the label (I). These structures represent various biphenyl and fluorene derivatives. The first two rows show biphenyls with substituents R<sub>1</sub>, R<sub>2</sub>, and groups Q and Z. The third row shows a fluorene derivative with substituents R<sub>1</sub>, R<sub>2</sub>, and groups Q and Z. The fourth row shows another fluorene derivative with substituents R<sub>1</sub>, R<sub>2</sub>, and groups Q and Z. The fifth row shows a biphenyl derivative with substituents R<sub>1</sub>, R<sub>2</sub>, and groups Q and Z. The sixth row shows a biphenyl derivative with substituents R<sub>1</sub>, R<sub>2</sub>, and groups Q and Z. The seventh row shows a biphenyl derivative with substituents R<sub>1</sub>, R<sub>2</sub>, and groups Q and Z. The eighth row shows a biphenyl derivative with substituents R<sub>1</sub>, R<sub>2</sub>, and groups Q and Z.


15  $Z$  is either,  $C \equiv C$ ,  $CR' = CR'$ ,  $CR'_2 - CR'_2$ ,  
 $C \equiv C - C \equiv C$ ,  $CR' = CR' - CR' = CR'$ ,  $C \equiv C - CR' = CR'$ , or  $CR'_2 -$   
 $CR'_2 - CR'_2 - CR'_2$  (where each  $R'$  independently represents H  
or a lower alkyl group);

(wherein each R'' independently is H, F, Cl, Br, I, a

-10-

- lower alkyl group,  $(\text{CH}_2)_n\text{OR}'$  where  $n=1, 2, \text{ or } 3$ ,  $\text{CF}_3$ ,  $\text{CH}_2\text{-CH}_2\text{F}$ ,  $\text{O-CH}_2\text{-CH}_2\text{F}$ ,  $\text{CH}_2\text{-CH}_2\text{-CH}_2\text{F}$ ,  $\text{O-CH}_2\text{-CH}_2\text{-CH}_2\text{F}$ ,  $\text{CN}$ ,  $(\text{C=O})\text{-R}'$ ,  $\text{N(R}')}_2$ ,  $\text{NO}_2$ ,  $(\text{C=O})\text{N(R}')}_2$ ,  $\text{O(CO)R}'$ ,  $\text{OR}'$ ,  $\text{SR}'$ ,  $\text{COOR}'$ ,  $\text{R}_{\text{Ph}}$ ,  $\text{CR}'=\text{CR}'\text{-R}_{\text{Ph}}$ ,  $\text{CR}'_2\text{-CR}'_2\text{-R}_{\text{Ph}}$  (where  $\text{R}_{\text{Ph}}$  represents an unsubstituted or substituted phenyl group with the phenyl substituents being chosen from any of the non-phenyl substituents defined for  $\text{R}''$ , a tri-alkyl tin, a tetrazole or oxadiazole of the form:

10




wherein  $\text{R}'$  is H or a lower alkyl group)

or  $\text{X}$  is  $\text{CR}'=\text{CR}'$ ,  $\text{N=N}$ ,  $\text{C=O}$ ,  $\text{O}$ ,  $\text{NR}'$  (where  $\text{R}'$  represents H or a lower alkyl group),  $\text{S}$ , or  $\text{SO}_2$ ;


- each  $\text{R}_1$  and  $\text{R}_2$  independently is H, F, Cl, Br, I, a lower alkyl group,  $(\text{CH}_2)_n\text{OR}'$  where  $n=1, 2, \text{ or } 3$ ,  $\text{CF}_3$ ,  $\text{CH}_2\text{-CH}_2\text{F}$ ,  $\text{O-CH}_2\text{-CH}_2\text{F}$ ,  $\text{CH}_2\text{-CH}_2\text{-CH}_2\text{F}$ ,  $\text{O-CH}_2\text{-CH}_2\text{-CH}_2\text{F}$ ,  $\text{CN}$ ,  $(\text{C=O})\text{-R}'$ ,  $\text{N(R}')}_2$ ,  $\text{NO}_2$ ,  $(\text{C=O})\text{N(R}')}_2$ ,  $\text{O(CO)R}'$ ,  $\text{OR}'$ ,  $\text{SR}'$ ,  $\text{COOR}'$ , a tri-alkyl tin,  $\text{R}_{\text{Ph}}$ ,  $\text{CR}'=\text{CR}'\text{-R}_{\text{Ph}}$ ,  $\text{CR}'_2\text{-CR}'_2\text{-R}_{\text{Ph}}$  (where  $\text{R}_{\text{Ph}}$  represents an unsubstituted or substituted phenyl group with the phenyl substituents being chosen from any of the non-phenyl substituents defined for  $\text{R}_1$  and  $\text{R}_2$ , a tetrazole or oxadiazole of the form:

25




(wherein  $\text{R}'$  is H or a lower alkyl group), or a triazene of the form:

-11-

(wherein  $R_8$  and  $R_9$  are lower alkyl groups)


or



5 each  $Q$  is independently selected from one of the following structures:

IA, IB, IC, ID, IE, IF and IG, wherein

IA has the following structure:



10


wherein:

each of  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$ , or  $R_7$  independently is defined the same as  $R_1$  above;


IB has the following structure:

(IB)

15



or




wherein:

each of  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$ ,  $R_{13}$ ,  $R_{14}$ ,  $R_{15}$ , or  $R_{16}$  independently

-12-

is defined the same as  $R_1$  above;


IC has the following structure:



wherein:

- 5 each of  $R_{17}$ ,  $R_{18}$ ,  $R_{19}$ ,  $R_{20}$ , or  $R_{21}$  independently is defined the same as  $R_1$  above;


ID has the following structure:



wherein:


- 10 each of  $R_{22}$ ,  $R_{23}$ , or  $R_{24}$  independently is defined the same as  $R_1$  above

and




represents a heterocyclic ring of one of the six following formulas:

-13-




IE has the following structure:




wherein:

- 5 each of  $R_{25}$ ,  $R_{26}$ , or  $R_{27}$  independently is defined the same as  $R_1$  above

and




represents a heterocyclic ring of one of the six following formulas:



-14-

IF has the following structure:




wherein:

exactly one of  $R_{28}$ ,  $R_{29}$ ,  $R_{30}$ ,  $R_{31}$  or  $R_{32}$  is the  $Z$


- 5 link defined for Formula I above and each other  $R_{28}$ ,  $R_{29}$ ,  $R_{30}$ ,  $R_{31}$  or  $R_{32}$  independently is defined the same as  $R_1$  above;

IG has the following structure:

(IG)



10




wherein:



-15-

exactly one of  $R_{33}$ ,  $R_{34}$ ,  $R_{35}$ ,  $R_{36}$ ,  $R_{37}$ ,  $R_{38}$ , or  $R_{39}$  is the

 link defined for Formula I above and each other

$R_{33}$ ,  $R_{34}$ ,  $R_{35}$ ,  $R_{36}$ ,  $R_{37}$ ,  $R_{38}$ , or  $R_{39}$  independently is defined the same as  $R_1$  above.

5 It is another object of the present invention to provide an amyloid binding compound of Formula I, as defined above, or a water soluble, non-toxic salt thereof, wherein at least one of the substituents  $R_1$ - $R_7$  and  $R_{10}$ - $R_{39}$  is selected from the group consisting of  $^{131}\text{I}$ ,  
10  $^{123}\text{I}$ ,  $^{76}\text{Br}$ ,  $^{75}\text{Br}$ ,  $^{18}\text{F}$ ,  $^{19}\text{F}$ ,  $^{125}\text{I}$ ,  $\text{CH}_2\text{-CH}_2\text{-}^{18}\text{F}$ ,  $\text{O-CH}_2\text{-CH}_2\text{-}^{18}\text{F}$ ,  $\text{CH}_2\text{-CH}_2\text{-}^{18}\text{F}$ ,  $\text{O-CH}_2\text{-CH}_2\text{-CH}_2\text{-}^{18}\text{F}$  and a carbon-containing substituent as specified in Formula I wherein at least one carbon is  $^{11}\text{C}$  or  $^{13}\text{C}$ .

15 It is a further object of the invention to provide an amyloid binding compound of Formula I, as defined above, or a water-soluble, non-toxic salt thereof, wherein the compound binds to  $\text{A}\beta$  with a dissociation constant ( $K_D$ ) between 0.0001 and 10.0  $\mu\text{M}$  when measured by binding to synthetic  $\text{A}\beta$  peptide or Alzheimer's Disease  
20 brain tissue.

Still another object of the present invention is to provide a method for synthesizing an amyloid binding compound of Formula I, as defined above, or a water soluble, non-toxic salt thereof, wherein at least one of  
25 the substituents  $R_1$ - $R_7$  and  $R_{10}$ - $R_{39}$  is selected from the group consisting of  $^{131}\text{I}$ ,  $^{123}\text{I}$ ,  $^{76}\text{Br}$ ,  $^{75}\text{Br}$ ,  $^{18}\text{F}$  and  $^{19}\text{F}$ , comprising the step of reacting an amyloid binding compound of Formula I, as defined above, or a water soluble, non-toxic salt thereof wherein at least one of  
30 the substituents  $R_1$ - $R_7$  and  $R_{10}$ - $R_{39}$  is a tri-alkyl tin, with a halogenating agent containing  $^{131}\text{I}$ ,  $^{123}\text{I}$ ,  $^{76}\text{Br}$ ,  $^{75}\text{Br}$ ,  $^{18}\text{F}$  or  $^{19}\text{F}$ .

An additional object of the present invention is a pharmaceutical composition for *in vivo* imaging of amyloid  
35 deposits, comprising (a) an amyloid binding compound of

-16-

Formula I, as defined above, or a water soluble, non-toxic salt thereof, wherein at least one of the substituents  $R_1$ - $R_7$  and  $R_{10}$ - $R_{39}$  is selected from the group consisting of  $^{131}\text{I}$ ,  $^{123}\text{I}$ ,  $^{76}\text{Br}$ ,  $^{75}\text{Br}$ ,  $^{18}\text{F}$ ,  $^{19}\text{F}$  and a carbon-containing substituent as specified in Formula I wherein at least one carbon is  $^{11}\text{C}$  or  $^{13}\text{C}$ , and (b) a pharmaceutically acceptable carrier.

Yet another object of the present invention is an *in vivo* method for detecting amyloid deposits in a subject, comprising the steps of: (a) administering a detectable quantity of the above pharmaceutical composition, and (b) detecting the binding of the compound to amyloid deposit in said subject. It is also an object of the present invention to provide an *in vivo* method for detecting amyloid deposits in a subject wherein the amyloid deposit is located in the brain of a subject. This method of the invention may be used in a subject who is suspected of having an amyloidosis associated disease or syndrome selected from the group consisting of Alzheimer's Disease, Down's Syndrome, and homozygotes for the apolipoprotein E4 allele.

Another object of the invention relates to pharmaceutical compositions and methods of preventing cell degeneration and toxicity associated with fibril formation in amyloidosis associated conditions such as AD and Type 2 diabetes mellitus. Such pharmaceutical compositions comprise alkyl, alkenyl and alkynyl derivatives of Chrysamine G and a pharmaceutically acceptable carrier. Such compounds would be non-toxic.

Another object of the invention relates to the use of the probes as a stain for the visualization and detection of amyloid deposits in biopsy or post-mortem tissue specimens.

Another object of this invention relates to the use of radiolabeled probes for the quantitation of amyloid deposits in biopsy or postmortem tissue specimens.

Another object relates to a method of distinguishing

-17-

Alzheimer's disease brain from normal brain.

Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

#### Brief Description of the Drawings

Figure 1A illustrates the chemical structures of Chrysamine G and several Chrysamine G analogues or derivatives which have been synthesized and tested, including the 3-iodo derivative (3-ICG), the 3-iodo dimethoxy derivative (3-ICG(OMe)<sub>2</sub>), the dimethyl ester derivative (CG(COOMe)<sub>2</sub>), the phenol derivative, salicylic acid (SA), the aniline derivative (1/2CG), Congo red, the 3,3'-diiodo derivative (3,3'-I<sub>2</sub>CG), the 3,3'-dibromo derivative (3,3'-Br<sub>2</sub>CG), the 3,3'-dichloro derivative (3,3'-Cl<sub>2</sub>CG), the 3-bromo derivative (3-BrCG), and the 5-fluorosalicylic acid derivative ((5-FSA)CG). The numbers in the figure refer to each compound's K<sub>i</sub> (in μM) for inhibition of [<sup>14</sup>C]Chrysamine G binding to the synthetic peptide, Aβ(10-43).

Figure 1B illustrates the chemical structures of several Chrysamine G analogues or derivatives which have been synthesized and tested, including the 3,3'-dicarboxylic acid derivative (3,3'-(COOH)<sub>2</sub>CG), the 2,2'-disulfonic acid derivative of Chrysamine G (2,2'-(SO<sub>3</sub>)<sub>2</sub>CG), the 3-bromo, 3-isopropylsalicylic acid derivative (3-Br-(3-iPrSA)CG), the 3-isopropylsalicylic acid derivative ((3-iPrSA)CG), the 2,4-diphenol derivative (2,4-Diphenol), the γ-resorcylic acid

-18-

derivative ((6-OHSA)CG), the 3,3',5,5'-tetramethylbenzidine derivative (3,3',5,5'-(CH<sub>3</sub>)<sub>4</sub>CG), the 3,3'-dimethyl derivative (3,3'-(CH<sub>3</sub>)<sub>2</sub>CG), the 2,2'-dimethyl derivative (2,2'-(CH<sub>3</sub>)<sub>2</sub>CG), the benzisoxazole derivative (CG Benzisoxazole), and the 3-carboxy alkyne derivative (3-(COOH)-C3C). The numbers in the figure refer to each compound's K<sub>i</sub> (in μM) for inhibition of [<sup>14</sup>C]Chrysamine G binding to the synthetic peptide, Aβ(10-43).

10        Figures 2A-2K illustrate the chemical structures of the alkenyl derivative of Chrysamine G and alkenyl tri-alkyl tin derivatives of analogues of Chrysamine G, in particular heterocyclic analogues. Note that these structures represent one-half of a molecule which is  
15        symmetric around the wavy bond shown in the upper right, except that the tri-alkyl tin moiety may only be on one side of the biphenyl group. The tri-alkyl tin derivatives are stable intermediate and immediate precursors for the preparation of high specific activity  
20        halogenated radioactive derivatives. The heterocyclic analogues represent alternative means of placing weakly acidic moieties in the same structural position as the moderately acidic carboxylic acid group of Chrysamine G. These tri-alkyl tin precursor compounds are shown in  
25        their protonated form, yet those of skill in the art recognize that their deprotonated forms and tautomers also are embraced by these drawings.

- 2A) alkenyl derivative of Chrysamine G.  
2B) alkenyl tri-alkyl tin derivative of Chrysamine G;  
30        2C) alkenyl tri-alkyl tin derivative of the 3-Hydroxy-1,2-benzisoxazole analogue;  
2D) alkenyl tri-alkyl tin derivative of the phthalimide or isoindole-1,3(2H)-dione analogue;  
2E) alkenyl tri-alkyl tin derivative of the  
35        phthalhydrazide or 2,3-benzodiazine-1,4(2H,3H)-dione

-19-

- analogue;
- 2F) alkenyl tri-alkyl tin derivative of the 2,3-benzoxazine-1,4(3H)-dione analogue;
- 2G) alkenyl tri-alkyl tin derivative of the (2H)1,3-benzoxazine-2,4(3H)-dione analogue;
- 5 2H) alkenyl tri-alkyl tin derivative of the (3H)2-benzazine-1,3(2H)-dione analogue;
- 2I) alkenyl tri-alkyl tin derivative of the 1,8-Naphthalimide analogue.
- 10 2J) alkenyl tri-alkyl tin derivative of the tetrazole analogue .
- 2K) alkenyl tri-alkyl tin derivative of the oxadiazole analogue.

Figure 3 Displacement curves of [<sup>14</sup>C]Chrysamine G binding to A $\beta$ (10-43) by several structural analogues of Chrysamine G. Abbreviations refer to those used in Figure 1. Figure 3A) Chrysamine G (open triangles); (5-FSA)CG (filled diamonds); 3,3'-(COOH)<sub>2</sub>CG (filled squares); 2,2'-(SO<sub>3</sub>)<sub>2</sub>CG (filled circles). Figure 3B) Chrysamine G (open triangles); Congo red (open circles); aniline derivative (open inverted triangles); phenol derivative (open squares); salicylic acid (X's). Curves which show increased binding at higher concentrations do so because of the formation of micelles. Bedaux, F. et al., *Pharm. Weekblad* 98: 189 (1963).

Figure 4A is a Scatchard plot of Chrysamine G binding to A $\beta$ (10-43). The curved line represents a nonlinear least-squares fit to a two, independent binding site model. The straight lines represent the individual components.

Figure 4B is a Scatchard analysis of [<sup>14</sup>C]CG binding to typical control (diamonds) and AD brain samples (squares). The dashed line has the same slope as the AD line and is meant to aid in the comparison with the

-20-

control slope. This AD brain sample had 48 NP/x200 magnification, a  $K_D$  of 0.35  $\mu\text{M}$ , and a  $B_{\text{max}}$  of 790 fmol/ $\mu\text{g}$  protein. The control had a  $K_D$  of 0.48  $\mu\text{M}$ , and a  $B_{\text{max}}$  of 614 fmol/ $\mu\text{g}$  protein.

5        Figure 5 is a graph illustrating the linearity of the binding assay with respect to peptide concentration. Approximately 0.9  $\mu\text{g}$  of  $\text{A}\beta(10-43)$  was used in the typical assay.

10        Figure 6A is a graph illustrating the time course of association of Chrysamine G and  $\text{A}\beta(10-43)$ .

Figure 6B is the graphic illustration of the determination of the association rate constant ( $k_1$ ).

Figure 6C is a graph of the time course of dissociation of Chrysamine G from  $\text{A}\beta(10-43)$ .

15        Figure 7 is a graphic representation of a molecular model of the interaction between Chrysamine G and  $\text{A}\beta$ .

Figure 8A is a graph illustrating the correlation between the amount of [ $^{14}\text{C}$ ]Chrysamine G bound and the number of neuritic plaques (NP) in AD brain samples.

20        Figure 8B is a graph illustrating the correlation between the amount of [ $^{14}\text{C}$ ]Chrysamine G bound and the number of neurofibrillary tangles (NFT) in AD brain samples. In both Figure 8A and 8B, the x-axis represents the average NP or NFT count per field at x200 magnification in Bielschowsky stained sections of either  
25        the superior/middle frontal (n=10) or superior temporal cortex (n=10). The filled symbols and heavy lines indicate brains without amyloid angiopathy, the open symbols and dashed lines indicate brains with amyloid  
30        angiopathy. The y-axis represents total, absolute

-21-

[<sup>14</sup>C]Chrysamine G binding (fmol/μg protein) in homogenates of brain samples adjacent to those used for staining. Approximately 75 μg protein and 150 nM [<sup>14</sup>C]Chrysamine G were used.

5        Figures 9A, 9B and 9C. The binding of Chrysamine G to various brain areas in samples of AD brain having more than 20 NPs/x200 magnification, referred to as "High Plaque AD Brains", is shown in figure 9A. The binding of Chrysamine G to brain areas in samples of AD brain having  
10 less than 20 NPs/x200 magnification, referred to as "Low Plaque AD Brains", is shown in Figure 9B. The data points represent the ratio of [<sup>14</sup>C]Chrysamine G binding in the designated brain area to [<sup>14</sup>C]Chrysamine G binding in the cerebellum (CB) of the same brain. Horizontal bars  
15 represent the mean and error bars represent the standard error for control (circles), and AD brain (diamonds in 9A and 9B). Brain areas include the frontal pole (FP), head of caudate (CAU), superior/ middle frontal (SMF), superior temporal (ST), inferior parietal (IP), and  
20 occipital (OC) cortex. Asterisks indicate significant differences compared to control (\*p<0.05; \*\*p<0.001). Two Down's syndrome brain samples are indicated in Figure 9C. The diamonds in 9C represent a brain from a 23 year old Down's syndrome patient not yet symptomatic with AD.  
25 The triangles in 9C represent a 51 year old Down's syndrome patient who had developed AD as do the vast majority of Down's syndrome patients by their 40's.

Figure 10 is a graph illustrating the tissue levels of Chrysamine G in mice injected with [<sup>14</sup>C]Chrysamine G in  
30 the lateral tail vein and sacrificed at the times indicated. The open symbols and thin lines represent absolute radioactivity in units of cpm/g tissue (left axis). The closed symbols and solid lines represent the ratio of brain radioactivity to that in kidney (top) or  
35 blood (middle). The ratios are plotted on the right

-22-

axis.

Figure 11. TOP: Section from the inferior temporal lobe of AD brain stained with 1,4-bis(2-(3-carboxy-4-hydroxyphenyl)ethen-1-yl)-benzene by the method of Stokes and Trickey, *J. Clin. Pathol.* 26: 241-242 (1973). Visible are a large number of neuritic plaques, a neurofibrillary tangle and frequent neuropil threads. Cerebrovascular amyloid also is intensely stained (not shown). The photomicrograph was obtained using fluorescence microscopy. BOTTOM: Section of transgenic mouse brain [Tg(HuAPP695.SWE)2576; Hsiao et al. *Science* 274: 99-102 (1996)] similarly stained with 1,4-bis(2-(3-carboxy-4-hydroxyphenyl)ethen-1-yl)-benzene showing an intensely stained plaque.

Figure 12. A bar graph showing the effect of increasing concentrations of A $\beta$ (25-35) in the presence and absence of Chrysamine G on the cellular redox activity of rat pheochromocytoma (PC-12) cells as measured by 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide, MTT, reduction. The reduction product of MTT absorbs at 560 nm which is plotted on the vertical axis. The effect of A $\beta$ (25-35) alone is shown in the filled bars and shows a dose dependent decrease in MTT reduction. Significant differences from control (no A $\beta$ (25-35), no Chrysamine G) are shown in white inside the filled bars. The protective effect of 20  $\mu$ M Chrysamine G is shown in the open bars. Significant differences between MTT reduction in the presence and absence of Chrysamine G are shown in black inside the open bars.

Figure 13. A bar graph showing the protective effect of increasing concentrations of Chrysamine G against the A $\beta$ (25-35)-induced reduction of cellular redox activity of rat pheochromocytoma (PC-12) cells as



-23-

measured by 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide, MTT, reduction. The reduction product of MTT absorbs at 560 nm which is plotted on the vertical axis. The effect of Chrysamine G in the absence of A $\beta$ (25-35) is shown in the filled bars. There was no significant difference between control (no A $\beta$ (25-35), no Chrysamine G) and any of the concentrations of Chrysamine G in the absence of A $\beta$ (25-35). MTT reduction in the presence of 1  $\mu$ M A $\beta$ (25-35) and increasing concentrations of Chrysamine G is shown in the open bars. Significant differences in MTT reduction between the presence and absence of A $\beta$ (25-35) at each concentration of Chrysamine G are shown in white inside the filled bars. Significant differences in MTT reduction between the A $\beta$ (25-35) control (no Chrysamine G) and A $\beta$ (25-35) plus increasing concentrations of Chrysamine G are shown in black inside the open bars.

Figure 14. Comparison of the effects of Chrysamine-G and the inactive phenol derivative on the toxicity induced by A $\beta$ (25-35). 1  $\mu$ M A $\beta$ (25-35) was present in all experiments except control. Chrysamine-G showed protective effects at 0.1 and 1  $\mu$ M, but the phenol derivative showed no protective effects, and perhaps enhanced the toxicity of A $\beta$ .

#### Detailed Description of the Preferred Embodiments

The present invention exploits the ability of alkyl, alkenyl and alkynyl Chrysamine G derivatives and radiolabeled derivatives thereof to cross the blood brain barrier *in vivo* and bind to A $\beta$  deposited in plaques, to A $\beta$  deposited in cerebrovascular amyloid, and to the amyloid consisting of the protein deposited in NFT. Chrysamine G is a Congo red derivative with the key structural difference being that the sulfonic acid moieties found in Congo red are replaced by carboxylic acid groups in Chrysamine G (Figure 1). This structural

-24-

alteration allows Chrysamine G to enter the brain better than Congo red and large macromolecules such as antibodies. Tubis et al., *J. Am. Pharmaceut. Assn.* 49: 422 (1960). Also, Chrysamine G may be a more specific  
5 marker for AD pathophysiology than antibodies, which would also label non-fibrillar A $\beta$  deposits of uncertain pathological significance.

The alkyl, alkenyl and alkynyl Chrysamine G derivatives of the present invention have each of the  
10 following characteristics: (1) specific binding to synthetic A $\beta$  *in vitro*, (2) binding to  $\beta$ -sheet fibril deposits in brain sections (3) ability to cross a non-compromised blood brain barrier *in vivo*.

The method of this invention determines the presence  
15 and location of amyloid deposits in an organ or body area, preferably brain, of a patient. The present method comprises administration of a detectable quantity of a pharmaceutical composition containing a compound of Formula I, as defined above, called a "detectable  
20 compound," or a pharmaceutically acceptable water-soluble salt thereof, to a patient. A "detectable quantity" means that the amount of the detectable compound that is administered is sufficient to enable detection of binding of the compound to amyloid. An "imaging effective  
25 quantity" means that the amount of the detectable compound that is administered is sufficient to enable imaging of binding of the compound to amyloid.

The invention employs amyloid probes which, in conjunction with non-invasive neuroimaging techniques  
30 such as magnetic resonance spectroscopy (MRS) or imaging (MRI), or gamma imaging such as positron emission tomography (PET) or single-photon emission computed tomography (SPECT), are used to quantify amyloid deposition *in vivo*. The term "*in vivo* imaging" refers  
35 to any method which permits the detection of a labeled compound of Formula I, as described above, of the present invention. For gamma imaging, the radiation emitted from

-25-

the organ or area being examined is measured and expressed either as total binding or as a ratio in which total binding in one tissue is normalized to (for example, divided by) the total binding in another tissue of the same subject during the same *in vivo* imaging procedure. Total binding *in vivo* is defined as the entire signal detected in a tissue by an *in vivo* imaging technique without the need for correction by a second injection of an identical quantity of labeled compound along with a large excess of unlabeled, but otherwise chemically identical compound. A "subject" is a mammal, preferably a human, and most preferably a human suspected of having dementia.

For purposes of *in vivo* imaging, the type of detection instrument available is a major factor in selecting a given label. For instance, radioactive isotopes and  $^{19}\text{F}$  are particularly suitable for *in vivo* imaging in the methods of the present invention. The type of instrument used will guide the selection of the radionuclide or stable isotope. For instance, the radionuclide chosen must have a type of decay detectable by a given type of instrument. Another consideration relates to the half-life of the radionuclide. The half-life should be long enough so that it is still detectable at the time of maximum uptake by the target, but short enough so that the host does not sustain deleterious radiation. The radiolabeled compounds of the invention can be detected using gamma imaging wherein emitted gamma irradiation of the appropriate wavelength is detected. Methods of gamma imaging include, but are not limited to, SPECT and PET. Preferably, for SPECT detection, the chosen radiolabel will lack a particulate emission, but will produce a large number of photons in a 140-200 keV range. For PET detection, the radiolabel will be a positron-emitting radionuclide such as  $^{18}\text{F}$  which will annihilate to form two 511 keV gamma rays which will be detected by the PET camera.

-26-

In the present invention, amyloid binding compounds/probes are made which are useful for *in vivo* imaging and quantification of amyloid deposition. These compounds are to be used in conjunction with non-invasive neuroimaging techniques such as magnetic resonance spectroscopy (MRS) or imaging (MRI), positron emission tomography (PET), and single-photon emission computed tomography (SPECT). In accordance with this invention, the alkyl, alkenyl and alkynyl Chrysamine G derivatives may be labeled with  $^{19}\text{F}$  or  $^{13}\text{C}$  for MRS/MRI by general organic chemistry techniques known to the art. See, e.g., March, J. "ADVANCED ORGANIC CHEMISTRY: REACTIONS, MECHANISMS, AND STRUCTURE (3rd Edition, 1985), the contents of which are hereby incorporated by reference.

The alkyl, alkenyl and alkynyl Chrysamine G derivatives also may be radiolabeled with  $^{18}\text{F}$ ,  $^{11}\text{C}$ ,  $^{75}\text{Br}$ , or  $^{76}\text{Br}$  for PET by techniques well known in the art and are described by Fowler, J. and Wolf, A. in POSITRON EMISSION TOMOGRAPHY AND AUTORADIOGRAPHY (Phelps, M., Mazziota, J., and Schelbert, H. eds.) 391-450 (Raven Press, NY 1986) the contents of which are hereby incorporated by reference. The alkyl, alkenyl and alkynyl Chrysamine G derivatives also may be radiolabeled with  $^{123}\text{I}$  for SPECT by any of several techniques known to the art. See, e.g., Kulkarni, *Int. J. Rad. Appl. & Inst. (Part B)* 18: 647 (1991), the contents of which are hereby incorporated by reference. In addition, the alkyl, alkenyl and alkynyl Chrysamine G derivatives may be labeled with any suitable radioactive iodine isotope, such as, but not limited to  $^{131}\text{I}$ ,  $^{125}\text{I}$ , or  $^{123}\text{I}$ , by iodination of a diazotized amino derivative directly via a diazonium iodide, see Greenbaum, F. *Am. J. Pharm.* 108: 17 (1936), or by conversion of the unstable diazotized amine to the stable triazene, or by conversion of a non-radioactive halogenated precursor to a stable tri-alkyl tin derivative which then can be converted to the iodo compound by several methods well known to the art. See,

-27-

Satyamurthy and Barrio *J. Org. Chem.* 48: 4394 (1983), Goodman et al., *J. Org. Chem.* 49: 2322 (1984), and Mathis et al., *J. Labell. Comp. and Radiopharm.* 1994: 905; Chumpradit et al., *J. Med. Chem.* 34: 877 (1991); Zhuang et al., *J. Med. Chem.* 37: 1406 (1994); Chumpradit et al., *J. Med. Chem.* 37: 4245 (1994). For example, a stable triazene or tri-alkyl tin derivative of alkyl, alkenyl and alkynyl Chrysamine G derivatives is reacted with a halogenating agent containing  $^{131}\text{I}$ ,  $^{125}\text{I}$ ,  $^{123}\text{I}$ ,  $^{76}\text{Br}$ ,  $^{75}\text{Br}$ ,  $^{18}\text{F}$  or  $^{19}\text{F}$ . Thus, the stable triazene and tri-alkyl tin derivatives of Chrysamine G and its analogues are novel precursors useful for the synthesis of many of the radiolabeled compounds within the present invention. As such, these triazene and tri-alkyl tin derivatives are one embodiment of this invention.

The alkyl, alkenyl and alkynyl Chrysamine G derivatives also may be radiolabeled with known metal radiolabels, such as Technetium-99m ( $^{99\text{m}}\text{Tc}$ ). Modification of the substituents to introduce ligands that bind such metal ions can be effected without undue experimentation by one of ordinary skill in the radiolabeling art. The metal radiolabeled alkyl, alkenyl and alkynyl Chrysamine G derivative can then be used to detect amyloid deposits.

The methods of the present invention may use isotopes detectable by nuclear magnetic resonance spectroscopy for purposes of *in vivo* imaging and spectroscopy. Elements particularly useful in magnetic resonance spectroscopy include  $^{19}\text{F}$  and  $^{13}\text{C}$ .

Suitable radioisotopes for purposes of this invention include beta-emitters, gamma-emitters, positron-emitters, and x-ray emitters. These radioisotopes include  $^{131}\text{I}$ ,  $^{123}\text{I}$ ,  $^{18}\text{F}$ ,  $^{11}\text{C}$ ,  $^{75}\text{Br}$ , and  $^{76}\text{Br}$ . Suitable stable isotopes for use in Magnetic Resonance Imaging (MRI) or Spectroscopy (MRS), according to this invention, include  $^{19}\text{F}$  and  $^{13}\text{C}$ . Suitable radioisotopes for *in vitro* quantification of amyloid in homogenates of biopsy or post-mortem tissue include  $^{125}\text{I}$ ,  $^{14}\text{C}$ , and  $^3\text{H}$ . The

-28-

preferred radiolabels are  $^{18}\text{F}$  for use in PET *in vivo* imaging,  $^{123}\text{I}$  for use in SPECT imaging,  $^{19}\text{F}$  for MRS/MRI, and  $^3\text{H}$  or  $^{14}\text{C}$  for *in vitro* studies. However, any conventional method for visualizing diagnostic probes can be utilized in accordance with this invention.

The method could be used to diagnose AD in mild or clinically confusing cases. This technique would also allow longitudinal studies of amyloid deposition in human populations at high risk for amyloid deposition such as Down's syndrome, familial AD, and homozygotes for the apolipoprotein E4 allele. Corder et al., *Science* 261: 921 (1993). A method that allows the temporal sequence of amyloid deposition to be followed can determine if deposition occurs long before dementia begins or if deposition is unrelated to dementia. This method can be used to monitor the effectiveness of therapies targeted at preventing amyloid deposition.

Generally, the dosage of the detectably labeled alkyl, alkenyl and alkynyl Chrysamine G derivative will vary depending on considerations such as age, condition, sex, and extent of disease in the patient, contraindications, if any, concomitant therapies and other variables, to be adjusted by a physician skilled in the art. Dosage can vary from 0.001 mg/kg to 1000 mg/kg, preferably 0.1 mg/kg to 100 mg/kg.

Administration to the subject may be local or systemic and accomplished intravenously, intraarterially, intrathecally (via the spinal fluid) or the like. Administration may also be intradermal or intracavitary, depending upon the body site under examination. After a sufficient time has elapsed for the compound to bind with the amyloid, for example 30 minutes to 48 hours, the area of the subject under investigation is examined by routine imaging techniques such as MRS/MRI, SPECT, planar scintillation imaging, PET, and emerging imaging techniques, as well. The exact protocol will necessarily vary depending upon factors specific to the patient, as

-29-

noted above, and depending upon the body site under examination, method of administration and type of label used; the determination of specific procedures would be routine to the skilled artisan. For brain imaging, preferably, the amount (total or specific binding) of the bound radioactively labelled Chrysamine G or Chrysamine G derivative or analogue is measured and compared (as a ratio) with the amount of labelled Chrysamine G or Chrysamine G derivative bound to the cerebellum of the patient. This ratio is then compared to the same ratio in age-matched normal brain.

The pharmaceutical compositions of the present invention are advantageously administered in the form of injectable compositions. A typical composition for such purpose comprises a pharmaceutically acceptable carrier. For instance, the composition may contain about 10 mg of human serum albumin and from about 0.5 to 500 micrograms of the labeled alkyl, alkenyl and alkynyl Chrysamine G derivative per milliliter of phosphate buffer containing NaCl. Other pharmaceutically acceptable carriers include aqueous solutions, non-toxic excipients, including salts, preservatives, buffers and the like, as described, for instance, in REMINGTON'S PHARMACEUTICAL SCIENCES, 15th Ed. Easton: Mack Publishing Co. pp. 1405-1412 and 1461-1487 (1975) and THE NATIONAL FORMULARY XIV., 14th Ed. Washington: American Pharmaceutical Association (1975), the contents of which are hereby incorporated by reference.

Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oil and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, saline solutions, parenteral vehicles such as sodium chloride, Ringer's dextrose, etc. Intravenous vehicles include fluid and nutrient replenishers. Preservatives include antimicrobials, anti-oxidants, chelating agents and inert gases. The pH and exact concentration of the various

-30-

components of the pharmaceutical composition are adjusted according to routine skills in the art. See, Goodman and Gilman's THE PHARMACOLOGICAL BASIS FOR THERAPEUTICS (7th Ed.).

5        Particularly preferred pharmaceutical compositions of the present invention are those that, in addition to specifically binding amyloid *in vivo* and capable of crossing the blood brain barrier, are also non-toxic at appropriate dosage levels and have a satisfactory  
10        duration of effect.

#### Molecular Modeling

      Molecular modeling was done on an Evans and Sutherland PS-330 computer graphics system, running the computer modeling program MacroModel (Version 2.5  
15        available from C. Still at Columbia University) to generate the A $\beta$  peptide chains in the anti-parallel beta-sheet conformation. Kirschner et al., *Proc. Natl. Acad. Sci. U.S.A.* 83: 503 (1986). The amyloid peptides were used without further structural refinement. The A $\beta$   
20        peptides were aligned so that alternate chains were spaced 4.76 Å apart, characteristic of beta-sheet fibrils. Kirschner, *supra*. Chrysamine G was energy minimized and aligned with the fibril model to maximize contact with lysine-16 of A $\beta$ (10-43) and the hydrophobic  
25        phenylalanine-19 and -20 region.

#### Characterization of Specific Binding to A $\beta$ Synthetic Peptide: Affinity, Kinetics, Maximum Binding

      The characteristics of Chrysamine G and Chrysamine G derivative binding is first analyzed using synthetic A $\beta$   
30        peptide called A $\beta$ (10-43). The 10-43 peptide was chosen because it has been shown that this peptide provides a model system containing all of the characteristic structural features of A $\beta$  peptides. Hilbich et al., *J. Mol. Biol.* 218: 149 (1991). The 10-43 amino acid  
35        fragment of A $\beta$  was synthesized with 9-fluorenylmethyl



-31-

chloroformate (Fmoc) chemistry by the Peptide Synthesis Facility of the University of Pittsburgh. The peptide was characterized by mass spectrometry and the major component had an  $M_r$  of 3600 g/mole (calc. 3598). The peptide was further purified by the method of Hilbich et al. which, in brief, consisted of sequential size-exclusion chromatography on a Biogel P10 column (2 x 180 cm, 200-400 mesh, Biorad, Richmond, CA) in 70% formic acid followed by a second elution through a Biogel P4 column (2 x 180 cm, 200-400 mesh) in 1M acetic acid. Hilbich et al., *J. Mol. Biol.* 218: 149 (1991). The peptide was lyophilized and stored at -80°C until used in the binding assays.

Amino acid sequence for A $\beta$ (10-43) is as follows:

|     |     |     |     |     |     |     |     |     |     |     |     |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|
| 10  | 11  | 12  | 13  | 14  | 15  | 16  | 17  | 18  | 19  | 20  | 21  |
| Tyr | Glu | Val | His | His | Gln | Lys | Leu | Val | Phe | Phe | Ala |
|     |     |     |     |     |     |     |     |     |     |     |     |
| 22  | 23  | 24  | 25  | 26  | 27  | 28  | 29  | 30  | 31  | 32  | 33  |
| Glu | Asp | Val | Gly | Ser | Asn | Lys | Gly | Ala | Ile | Ile | Gly |
|     |     |     |     |     |     |     |     |     |     |     |     |
| 34  | 35  | 36  | 37  | 38  | 39  | 40  | 41  | 42  | 43  |     |     |
| Leu | Met | Val | Gly | Gly | Val | Val | Ile | Ala | Thr |     |     |

#### Binding assay to synthetic A $\beta$ (10-43)

Binding assays were performed in 12 x 75 mm borosilicate glass tubes. Various concentrations of nonradioactive Chrysamine G derivatives were added in 10% ethanol/water. Ethanol was necessary to prevent the micelle formation which occurs with these diazo dye derivatives, since the micelles are trapped by the filter even in the absence of peptide. To the above solution, 25  $\mu$ l of a 0.36 mg/ml suspension of A $\beta$ (10-43) in H<sub>2</sub>O was added and 10% ethanol was added to bring the volume to 950  $\mu$ l. After incubating for 10 min at room temperature,

-32-

50  $\mu$ l of [ $^{14}$ C]Chrysamine G in 40% ethanol was added, resulting in a final concentration of [ $^{14}$ C]Chrysamine G of 100-125 nM depending on the preparation of [ $^{14}$ C]Chrysamine G used. The binding mixture was incubated for 30 min at room temperature. Bound and free radioactivity were separated by vacuum filtration through Whatman GF/B filters using a Brandel M-24R Cell Harvester (Gaithersburg, MD) followed by two 3-ml washes with 10% ethanol at room temperature. Filters were equilibrated overnight in 4 ml Cytoscint®-ES scintillant (ICN Biomedicals, Inc., Irvine, CA) in 7.0 ml plastic scintillation vials before counting. In this and all binding assays, incubations were done at least in triplicate and the results expressed as mean  $\pm$  standard deviation.

#### Kinetic studies

Kinetics studies of [ $^{14}$ C]Chrysamine G binding to A $\beta$ (10-43) were performed in 13 x 100 mm borosilicate glass tubes by the filtration assay described above. For the kinetics of association, 25  $\mu$ l of 0.36 mg/ml A $\beta$ (10-43) were placed in 475  $\mu$ l of 10% ethanol and 4.5 ml of 125 nM [ $^{14}$ C]Chrysamine G was added to the solution at time zero. The mixture was rapidly vortexed and the binding reaction was stopped by vacuum filtration through Whatman GF/B filters using a Brandel M-24R Cell Harvester (Gaithersburg, MD) followed by two 3-ml washes with 10% ethanol at room temperature at times of 5, 10, 20, 30, 45, 60, 75, 135, 240, and 300 sec.; bound radioactivity was determined as above.

For the kinetics of dissociation, 25  $\mu$ l of 0.36 mg/ml A $\beta$ (10-43) were placed in 450  $\mu$ l of 10% ethanol followed by 25  $\mu$ l of 2.5  $\mu$ M [ $^{14}$ C]Chrysamine G in 40% ethanol. This mixture was vortexed and incubated for 30 min at room temperature. The mixture was diluted with 4.5 ml of 10  $\mu$ M nonradioactive Chrysamine G in 10% ethanol at time zero, the mixture was rapidly vortexed,

-33-

and the dissociation was stopped by filtration as above at times of 0.5, 1.5, 3, 5, and 15 min, and bound radioactivity was determined as above.

5     **Characterization of Specific Binding to Alzheimer's Disease Brain**

Binding of Chrysamine G to AD and control brain homogenates

          Autopsy brain samples were obtained from the Neuropathology Core of the Alzheimer's Disease Research  
10     Center of the University of Pittsburgh. Controls were defined as not meeting neuropathological criteria for AD (sufficient number of NPs or NFTs) according to the standards specified in a published NIA conference report. Khachaturian, *Arch. Neurol.* 42: 1097 (1985). Brain  
15     samples from eight control (ages 58-75), eleven AD (ages 61-84), and two Down syndrome brains (ages 23 and 51) were studied. There were six high-plaque (>20 NPs/x200 magnification) and five low-plaque (<20 NPs/x200 magnification) AD brains. Two controls were clinically  
20     demented but had no NPs or NFTs and received the diagnosis of "Dementia Lacking Distinctive Histology". Knopman *Dementia* 4: 132 (1993). Another control had dementia and olivopontocerebellar atrophy. The other controls had no clinical or histological evidence of  
25     neurologic disease. Autopsy samples were immediately frozen at -70 °C and stored at that temperature until homogenized. The numbers of NPs and NFTs were counted in sections of five separate but adjacent fields (x200 magnification) between cortical layers 2 and 4 in the  
30     cortex at the junction of superior and middle frontal gyri and superior temporal isocortex of all brains studied. A qualitative assessment of the presence of amyloid angiopathy in the superior/middle frontal cortex was made. The Bielschowsky silver impregnation method  
35     was used to identify NPs and NFTs and Congo red staining was used to identify cerebral amyloid angiopathy.

-34-

Details of this procedure have been previously published. Moossy et al., Arch. Neurol. 45: 251 (1988). Samples used for CG binding to the superior/middle frontal or superior temporal cortex were adjacent on gross  
5 dissection to those used for NP and NFT counts.

Approximately 100 mg of tissue from the junction of the superior and middle frontal cortex, superior temporal cortex, frontal pole, head of the caudate, inferior parietal cortex, occipital cortex, or cerebellum were  
10 homogenized with a Polytron® tissue homogenizer (PT 10/35, Brinkman Instruments Inc., Westbury, NY) for 30 sec at setting 6 in 10% ethanol at a concentration of 10-20 mg brain/ml. Not all areas were available from each brain. Aliquots of 25-150  $\mu$ l tissue (about 25-300  $\mu$ g of  
15 protein by the method of Lowry et al., J. Biol. Chem. 193: 265 (1951)) were incubated in 12 x 75 mm borosilicate glass tubes at room temperature with 10-750 nM [ $^{14}$ C]CG (26.8 Ci/mole) in a final volume of 1.0 ml of 10% ethanol for 30 min at room temperature. The standard  
20 conditions employed about 150  $\mu$ g of protein and 75 nM [ $^{14}$ C]CG for the cerebellar ratio studies and about 75  $\mu$ g of protein and 150 nM [ $^{14}$ C]CG for the correlative studies with NPs, NFTs, and amyloid angiopathy. Ethanol was  
25 necessary to prevent the micelle formation which occurs with diazo dye derivatives, since the micelles are trapped by the filter even in the absence of tissue. Bound and free radioactivity were separated by vacuum filtration through Whatman GF/B filters using a Brandel  
M-24R Cell Harvester (Gaithersburg, MD) followed by two  
30 3-ml washes with 10% ethanol at room temperature. Filters were equilibrated overnight in 4 ml Cytoscint®-ES scintillant (ICN Biomedicals, Inc., Irvine, CA) in 7.0 ml plastic scintillation vials before counting. Saturable (specific) binding was defined as total binding  
35 minus residual (non-saturable) binding in the presence of 20  $\mu$ M unlabelled CG. In all binding assays, incubations were done at least in triplicate and the results

-35-

expressed as mean  $\pm$  standard error unless otherwise specified. Results were expressed either in absolute terms of fmol [ $^{14}\text{C}$ ]CG bound/ $\mu\text{g}$  protein in a given brain area, or as a ratio of the fmol/ $\mu\text{g}$  protein in that brain area to the fmol/ $\mu\text{g}$  protein in the cerebellum of the same brain.

#### Octanol/Water Partitioning

Approximately 75  $\mu\text{M}$  solutions of Chrysamine G or its analogues were prepared in 5.0 ml 1-octanol. Five ml of phosphate buffered saline (0.15 M NaCl, 5 mM potassium phosphate, pH 7.4) were added and the layers mixed by rapid vortexing. The mixture was then centrifuged at 1,000 g to facilitate the formation of two clear phases. The layers were separated using a separatory funnel and 600  $\mu\text{l}$  of each layer was diluted with 400  $\mu\text{l}$  of ethanol and the absorbance measured at 389 nm for Chrysamine G or the  $\lambda_{\text{max}}$  for each analogue. Concentrations were determined after correction for the molar absorptivity differences in the two solvents and the partition coefficient expressed as the concentration in the octanol layer divided by the concentration in the aqueous layer. Experiments were done in triplicate.

#### Imaging the Binding of Chrysamine G to Amyloid Deposits in Alzheimer's Disease Brain

For visual demonstration of CG derivative binding to tissue, 8 micron paraffin sections of an AD brain with heavy deposits of cerebrovascular amyloid were stained with 1,4-bis(2-(3-carboxy-4-hydroxyphenyl)ethen-1-yl)-benzene by a modification of the method of Stokes and Trickey, *J. Clin. Pathol.* 26: 241-242 (1973) with 1 mM 1,4-bis(2-(3-carboxy-4-hydroxyphenyl)ethen-1-yl)-benzene substituted for Congo red, but the procedure was otherwise identical. Stained slides were examined using fluorescence microscopy.

-36-

**Determining Compound's Ability to Cross  
the Blood Brain Barrier**

Mouse studies Female Swiss-Webster mice were injected in the lateral tail vein with approximately 0.03  $\mu\text{Ci/g}$  of [ $^{14}\text{C}$ ]Chrysamine G in a 0.9% NaCl solution. Mice were sacrificed by cervical dislocation at intervals of 15 min, 35 min, 1 hr, 4 hr, and 24 hr after injection. The carotid blood, brain, liver, and kidneys were rapidly obtained, weighed, and homogenized in distilled/deionized  $\text{H}_2\text{O}$  using a ground glass homogenizer. An aliquot was weighed into an 18.0 ml plastic scintillation vial (Beckman Poly-Q-Vial) and counted after addition of 10.0 ml of scintillation cocktail (Cytoscint®-ES (ICN)) and overnight equilibration. The [ $^{14}\text{C}$ ]Chrysamine G content of the tissues was expressed as cpm/mg tissue.

Experiments in which radioactivity was extracted from tissues were performed as above except 0.05  $\mu\text{Ci/g}$  of [ $^{14}\text{C}$ ]Chrysamine G was injected and the mice were sacrificed at 60 min. Brain and liver were then removed and extracted with a Folch procedure. Folch et al., *J. Biol. Chem.* 226: 447 (1957). In both tissues, over 95% of the extracted radioactivity was contained in the organic layer. The organic layer was evaporated to dryness, resuspended in a minimal amount of 10% methanol/90% ACN, and injected onto a silica column (Prep Nova Pak HR Silica, 7.8 x 300 mm, Waters, Milford, MA) and eluted isocratically with the same solvent. Under these conditions, 99% of the radioactivity is eluted in the solvent front, but most lipids are retained longer, making the fraction eluting in the solvent front suitable for injection onto the reverse-phase C4 column system described above. The entire solvent front was collected, dried, and resuspended in 10% ACN/90% sodium phosphate buffer (5mM, pH 6) and injected, along with authentic non-radioactive Chrysamine G, onto the C4 column. One minute fractions were collected and counted after addition of 10 ml of Cytoscint®-ES.

-37-

In yet another embodiment, the invention relates to a pharmaceutical composition and method for preventing cell degeneration and toxicity associated with fibril formation in certain "amyloidosis associated" conditions

5 such as Alzheimer's Disease, Down's Syndrome and Type 2 diabetes mellitus, hereditary cerebral hemorrhage amyloidosis (Dutch), amyloid A (reactive), secondary amyloidosis, familial mediterranean fever, familial amyloid nephropathy with urticaria and deafness

10 (Muckle-wells Syndrome), amyloid lambda L-chain or amyloid kappa L-chain (idiopathic, myeloma or macroglobulinemia-associated) A beta 2M (chronic hemodialysis), ATTR (familial amyloid polyneuropathy (Portuguese, Japanese, Swedish), familial amyloid

15 cardiomyopathy (Danish), isolated cardiac amyloid, (systemic senile amyloidosises), AIAPP or amylin insulinoma, atrial natriuretic factor (isolated atrial amyloid), procalcitonin (medullary carcinoma of the thyroid), gelsolin (familial amyloidosis (Finnish)),

20 cystatin C (hereditary cerebral hemorrhage with amyloidosis (Icelandic)), AApo-A-I (familial amyloidotic polyneuropathy - Iowa), AApo-A-II (accelerated senescence in mice), fibrinogen-associated amyloid; and Asor or Pr P-27 (scrapie, Creutzfeldt Jacob disease,

25 Gertsman-Straussler-Scheinker syndrome, bovine spongiform encephalitis) or in cases of persons who are homozygous for the apolipoprotein E4 allele. This method involves administering a pharmaceutical composition comprising Chrysamine G, or one of the above described

30 derivatives thereof, to a subject suspected of having or at high risk of developing such amyloidosis associated condition.

Because certain diazo compounds could be carcinogenic, the therapeutic compounds of the present

35 invention include only non-toxic, non-carcinogenic compounds. That is, the present invention addresses the problems with potential carcinogenicity by using only

-38-

alkyl, alkenyl and alkynyl compounds which do not contain groups that could be metabolized to carcinogenic benzidine compounds.

Any potential problems with lower bioavailability also is avoided by the use of alkyl, alkenyl and alkynyl derivatives of the azo compounds. These compounds are not substrates for reduction by bacterial or mammalian azo reductases.

Indeed, compounds of the present invention intended for therapeutic use are advantageous over existing compounds because they contain an 1,4-bis(2-(3-carboxy-4-hydroxyphenyl)ethen-1-yl)-benzene linkage which is not a substrate for bacterial azo-reductases in the intestines.

*In vitro* studies have shown that A $\beta$  neurotoxicity requires fibril formation and is inhibited by Congo red. Specifically, it has been shown that the amyloid fibril-binding dye Congo red inhibits fibrillar A $\beta$  neurotoxicity by inhibiting fibril formation or by binding preformed fibrils. Lorenzo et al., *Proc. Natl. Acad. Sci. USA* 91: 12243-12247 (1994). Congo red also has been shown to inhibit pancreatic islet cell toxicity of diabetes-associated amylin, another type of amyloid fibril. Lorenzo et al., *supra*. See also, Burgevin et al. *NeuroReport* 5: 2429 (1994); Pollack et al., *J. Neurosci. Letters* 184: 113-116 (1995); Pollack et al. *Neuroscience Letters* 197: 211 (1995). These data indicate that amyloid-binding compounds such as alkyl, alkenyl and alkynyl derivatives of Chrysamine G, which are similar to Congo red but which, unlike Congo red, enter the brain well, would be effective in preventing cell degeneration and toxicity associated with fibril formation in amyloidosis associated conditions.

In Example 8 and Figures 12 and 13, it is shown that Chrysamine G has effects very similar to those previously reported for Congo red in having a dose-dependent, protective effect in rat pheochromocytoma. Therefore, these *in vitro* assays provide a means for selecting



-39-

compounds for use in pharmaceutical compositions for the prevention of cell degeneration and toxicity associated with fibril formation.

Compounds such as Chrysamine G and the above  
5 described derivatives thereof, are tested pursuant to the present invention, for *in vivo* efficacy in preventing amyloid fibril formation or associated cellular degeneration, as measured by the formation of dystrophic neurites, synapse loss, neurofibrillary tangle formation  
10 and gliosis, in an animal model, such as the "senile animal" model for cerebral amyloidosis, Wisniewski *et al.*, *J. Neuropathol. & Exp. Neurol.* 32: 566 (1973), the mouse model of familial Mediterranean fever (Neurochem., Inc. Kingston, Ontario, Canada) and the transgenic mouse  
15 model of Alzheimer-type neuropathology, Games *et al.*, *Nature* 373: 523-527 (1995); Hsiao *et al.* *Science* 274: 99-102 (1996). In the familial Mediterranean fever model, the animals develop systemic amyloidosis. In an  
20 *in vivo* assay according to this invention, serial necropsies in animals treated and untreated with the compounds of the present invention to evaluate the inhibition of amyloid formation are compared. In the animal models for cerebral amyloid formation, in addition to following amyloid formation serially, the presence of  
25 amyloid-associated neurodegeneration, as measured by the formation of dystrophic neurites, synapse loss, neurofibrillary tangle formation and gliosis, also is assessed in serial necropsies in animals treated and untreated with the compounds of the present invention.

30 According to the present invention, a pharmaceutical composition comprising Chrysamine G or derivatives thereof, is administered to subjects in whom amyloid or amyloid fibril formation, cell degeneration and toxicity are anticipated. In the preferred embodiment, such  
35 subject is a human and includes, for instance, those who are at risk of developing cerebral amyloid, including the elderly, nondemented population and patients having

-40-

amyloidosis associated diseases and Type 2 diabetes mellitus. The term "preventing" is intended to include the amelioration of cell degeneration and toxicity associated with fibril formation. By "amelioration" is meant the prevention of more severe forms of cell degeneration and toxicity in patients already manifesting signs of toxicity, such as dementia.

The pharmaceutical composition for purposes of preventing cell degeneration and toxicity associated with fibril formation in amyloidosis associated diseases comprises Chrysamine G or a derivative thereof described above and a pharmaceutically acceptable carrier. In one embodiment, such pharmaceutical composition comprises serum albumin, Chrysamine G or Chrysamine G derivative and a phosphate buffer containing NaCl. Other pharmaceutically acceptable carriers include aqueous solutions, non-toxic excipients, including salts, preservatives, buffers and the like, as described, for instance, in REMINGTON'S PHARMACEUTICAL SCIENCES, 15th Ed., Easton: Mack Publishing Co., pp. 1405-1412 and 1461-1487 (1975) and THE NATIONAL FORMULARY XIV., 14th Ed. Washington: American Pharmaceutical Association (1975), and the UNITED STATES PHARMACOPEIA XVIII. 18th Ed. Washington: American Pharmaceutical Association (1995), the contents of which are hereby incorporated by reference.

Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oil and injectable organic esters such as ethyl oleate. Aqueous carriers include water, alcoholic/aqueous solutions, saline solutions, parenteral vehicles such as sodium chloride, Ringer's dextrose, etc. Intravenous vehicles include fluid and nutrient replenishers. Preservatives include antimicrobial, anti-oxidants, chelating agents and inert gases. The pH and exact concentration of the various components the pharmaceutical composition are adjusted according to routine skills in the art. See, Goodman and

-41-

Gilman's THE PHARMACOLOGICAL BASIS FOR THERAPEUTICS (7th Ed.).

According to the invention, such pharmaceutical composition could be administered orally, in the form of  
5 a liquid or solid, or injected intravenously or intramuscularly, in the form of a suspension or solution. By the term "pharmaceutically effective amount" is meant an amount that prevents cell degeneration and toxicity associated with fibril formation. Such amount would  
10 necessarily vary depending upon the age, weight and condition of the patient and would be adjusted by those of ordinary skill in the art according to well-known protocols. In one embodiment, a dosage would be between 0.1 and 100 mg/kg per day, or divided into smaller  
15 dosages to be administered two to four times per day. Such a regimen would be continued on a daily basis for the life of the patient. Alternatively, the pharmaceutical composition could be administered intramuscularly in doses of .1 to 100 mg/kg every one to  
20 six weeks.

In yet another embodiment, the invention relates to a method of detecting amyloid deposits in biopsy or post-mortem tissue. The method involves incubating formalin-fixed tissue with a solution of a compound of Formula I,  
25 described above. Preferably, the solution is 25-100% ethanol, (with the remainder being water) saturated with the compound of Formula I. Upon incubation, the compound stains or labels the amyloid deposit in the tissue, and the stained or labelled deposit can be detected or  
30 visualized by any standard method. Such detection means include microscopic techniques such as bright-field, fluorescence, laser-confocal and cross-polarization microscopy.

In yet another embodiment, the invention relates to  
35 a method of quantifying the amount of amyloid in biopsy or post-mortem tissue. This method involves incubating a labelled alkyl, alkenyl and alkynyl derivative of

-42-

Chrysamine G, preferably the compounds of Formula I, or a water-soluble, non-toxic salt thereof, with homogenate of biopsy or post-mortem tissue. The tissue is obtained and homogenized by methods well known in the art. The preferred label is a radiolabel, although other labels such as enzymes, chemiluminescent and immunofluorescent compounds are well known to skilled artisans. The preferred radiolabel is  $^{125}\text{I}$ ,  $^{14}\text{C}$  or  $^3\text{H}$ , the preferred label substituent of Formula I is at least one of  $\text{R}_1\text{-R}_7$ ,  $\text{R}_{10}\text{-R}_{27}$ .

10 Tissue containing amyloid deposits will bind to the labeled alkyl, alkenyl and alkynyl derivatives of Chrysamine G. The bound tissue is then separated from the unbound tissue by any mechanism known to the skilled artisan, such as filtering. The bound tissue can then be

15 quantified through any means known to the skilled artisan. See Example 3. The units of tissue-bound radiolabeled Chrysamine G derivative are then converted to units of micrograms of amyloid per 100 mg of tissue by comparison to a standard curve generated by incubating

20 known amounts of amyloid with the radiolabeled Chrysamine G derivative.

In yet another embodiment, the invention relates to a method of distinguishing an Alzheimer's diseased brain from a normal brain involving obtaining tissue from (i)

25 the cerebellum and (ii) another area of the same brain, other than the cerebellum, from normal subjects and from subjects suspected of having Alzheimer's disease. See Example 3. Such tissues are made into separate homogenates using methods well known to the skilled

30 artisan, and then are incubated with a radiolabeled alkyl, alkenyl and alkynyl Chrysamine G derivative. The amount of tissue which binds to the radiolabeled alkyl, alkenyl and alkynyl Chrysamine G derivative is then calculated for each tissue type (e.g. cerebellum, non-

35 cerebellum, normal, abnormal) and the ratio for the binding of non-cerebellum to cerebellum tissue is calculated for tissue from normal and for tissue from

-43-

patients suspected of having Alzheimer's disease. These ratios are then compared. If the ratio from the brain suspected of having Alzheimer's disease is above 90% of the ratios obtained from normal brains, the diagnosis of Alzheimer's disease is made.

**EXAMPLE 1.     THE SYNTHESIS OF CHRYSAMINE G AND DERIVATIVES THEREOF**

**Synthesis of Chrysamine G**

The synthesis of Chrysamine G (i.e., 4,4'-bis(3-carboxy-4-hydroxyphenylazo)-biphenyl) requires the following reaction steps. These reaction steps will be referred to as the "Chrysamine G Synthesis" general procedure. Benzidine•2HCl (28.9 mg, 0.11 mmole, Sigma Chemical Company, St. Louis, MO) was added to 1.5 ml of 1:1 DMSO:distilled/deionized H<sub>2</sub>O in a 50cc round bottom flask. Each of the reaction steps were carried out at 0°C unless otherwise specified. Twenty-nine µl of concentrated HCl were added, resulting in a clear solution after stirring. To the benzidine solution, a solution of 15.5 mg (0.22 mmole) of NaNO<sub>2</sub> in 300 µl of 1:1 DMSO/H<sub>2</sub>O was added drop-wise, resulting in a pH of about 2-3. The reaction mixture was stirred for 45 min, and then to this tetra-azotized benzidine mixture was added drop-wise over a 10 min period to 24.8 mg (0.18 mmole) of methyl salicylate (Aldrich) dissolved in 2.0 ml of 100% DMSO containing 250 mg/ml Na<sub>2</sub>CO<sub>3</sub> in suspension, keeping the pH about 10.5. The resulting mixture was stirred for 1 hr at 0°C, and then overnight at room temperature.

After this time, the pH was adjusted to about 7 and the mixture was extracted with three 50 ml portions of chloroform. The combined chloroform extracts were washed with three 50 ml portions of H<sub>2</sub>O, and then taken to dryness yielding the dimethyl ester of Chrysamine G (i.e., 4,4'-bis(3-methoxycarbonyl-4-hydroxyphenylazo)-biphenyl), which was further purified by recrystallization from chloroform/hexane. The ester was

-44-

then hydrolysed by dissolution in about 100 ml of 1:1 ethanol:H<sub>2</sub>O containing four equivalents of NaOH and refluxed for three hours. Evaporation of the ethanol followed by lyophilization of the H<sub>2</sub>O yielded the tetra-sodium salt of Chrysamine G. The free acid of Chrysamine G was formed by dissolving the tetra-sodium salt in H<sub>2</sub>O, washing once with chloroform to remove any unhydrolysed dimethyl ester, lowering the pH to about 2 and extracting with three 50 ml portions of ethyl acetate. The combined ethyl acetate extracts were washed with three 50 ml portions of H<sub>2</sub>O and taken to dryness.

Under these conditions, there was no remaining methyl salicylate, salicylic acid, or benzidine, and only trace amounts of the mono-substituted product, 4-hydroxy-4'-(3-carboxy-4-hydroxyphenylazo)-biphenyl by reverse-phase HPLC using a C4 column (Vydac 214-TP510) using a solvent system of sodium phosphate buffer (5 mM, pH 6):acetonitrile (ACN) 90:10, isocratically, for 10 min and then increased to 50% ACN over the next 20 min at a flow rate of 3.5 ml/min. The column eluant was monitored at 290 and 365 nm with a dual wavelength, diode array detector (Perkin Elmer 235C). Under these conditions, Chrysamine G eluted at 17.6 min.

The structure of Chrysamine G and derivatives was confirmed by proton NMR at 500 MHz in DMSO-d<sub>6</sub> with TMS as the internal standard. The peak assignments for the tetra-sodium salt of Chrysamine G were as follows with SA referring to protons at the specified ring position on the salicylic acid moiety and BZ referring to protons on the benzidine moiety: SA-3, doublet J=8.73 Hz at 6.75 parts per million (ppm); SA-4, doublet of doublets J=8.73 and 2.72 Hz at 7.82 ppm; BZ-2/6, doublet J=8.44 Hz at 7.91 ppm; BZ-3/5, doublet J=8.44 Hz at 7.95 ppm; and SA-6, doublet J=2.72 Hz at 8.28 ppm. The UV/visible spectrum in 40% ethanol showed a  $\lambda_{\max}$  at 389 nm. The molar absorptivity of Chrysamine G was determined by calculating the concentration of Chrysamine G through

-45-

comparison of peak areas to an internal standard by NMR and then immediately running the UV/vis spectrum of an aliquot of the NMR sample diluted in 40% ethanol. The molar absorptivity in 40% ethanol at 389 nm was  $5.5 \times 10^4$  AU/(cm•M).

5        $[^{14}\text{C}]$ Chrysamine G was synthesized by a modification of the above procedure. The tetra-azotization of benzidine was performed as described above except in 100%  $\text{H}_2\text{O}$ . Fifty  $\mu\text{l}$  of 2.5 M  $\text{Na}_2\text{CO}_3$  in  $\text{H}_2\text{O}$  were added to 50  $\mu\text{Ci}$  of crystalline salicylic acid-carboxy- $^{14}\text{C}$  (Sigma) in a 0.5 ml conical glass vial. Sixty  $\mu\text{l}$  of the tetra-azotized benzidine mixture was added to the conical vial, vortexed and kept at  $0^\circ\text{C}$  for 1 hr. To prevent formation of the mono-substituted benzidine by-product, 12.5  $\mu\text{l}$  of 250 mM non-radioactive salicylic acid (Sigma) in 2.5 M  $\text{Na}_2\text{CO}_3$  was added to the reaction mixture and maintained for 1 hr at  $0^\circ\text{C}$ . The vial was kept overnight at room temperature. The entire mixture was dissolved in a minimal amount of 35% ACN and injected onto the C4 column as described above. The peak corresponding to the Chrysamine G standard was collected and lyophilized. A specific activity of 26.8 Ci/mole was calculated by determining the absorbance at 389 nm and then counting the radioactivity in an aliquot of the same sample. The  $[^{14}\text{C}]$ Chrysamine G was stored in 40% ethanol. When the purified  $[^{14}\text{C}]$ Chrysamine G was re-injected onto the C4 column and eluted isocratically with 21% ACN at 3.5 ml/min, > 98% of the radioactivity co-eluted with authentic Chrysamine G at 10.4 min. Many of the Chrysamine G derivatives were synthesized using this "Chrysamine G Synthesis" general procedure, with the exceptions noted below. Structures of the derivatives were verified by NMR. Figure 1 shows the chemical structure of Chrysamine G and several derivatives.

35       Likewise,  $[^3\text{H}]$ Chrysamine G was synthesized by a modification of the above procedure. 3,3'- $[^3\text{H}]$ Benzidine was commercially prepared by American Radiolabelled

-46-

Chemicals, Inc. (St. Louis) by treating a mixture of 0.7 mg of 3,3'-diiodobenzidine, 1 mg palladium/charcoal catalyst, 200  $\mu$ L N,N-diisopropylethylamine, and 500  $\mu$ L THF with 8.26 Ci of carrier-free tritium gas at room temperature for 3 hrs. Exact yield and specific activity were not provided. The 3,3'-[ $^3$ H]benzidine co-eluted with authentic benzidine and showed no mono-iodo impurities. Because the material commercially provided had some residual N,N-diisopropylethylamine which interfered with diazotization, the 3,3'-[ $^3$ H]benzidine was purified by elution from a Vydac C4 column with a linear gradient of 100% water (pH= 6.0) to 100% 0.01 N HCl (pH=2.2) in 10 minutes (4.0 ml/min). By this procedure, the 3,3'-[ $^3$ H]benzidine eluted at approximately 9.5 minutes free of N,N-diisopropylethylamine and in a solvent which could be used without modification in the diazotization reaction. Diazotization in a volume of several hundred  $\mu$ L with a 100-fold excess of sodium nitrite, followed by coupling to a 100-fold excess of salicylic acid in NaCO<sub>3</sub> resulted in the tritiated derivative of Chrysamine G (4,4'-bis(3-carboxy-4-hydroxyphenylazo)-3,3'-[ $^3$ H]biphenyl) which was purified as above by HPLC giving a specific activity of ~35 Ci/mmol.

#### 25      Synthesis of Alkenyl (CH=CH) derivatives of Chrysamine-G

4,4'-biphenyldicarboxylic acid (Aldrich) is converted by reduction with LiAlH<sub>4</sub> to 4,4'-bis(hydroxymethyl)biphenyl, which, in turn, is converted to 4,4'-bis(iodomethyl)biphenyl by reaction with NaI and BF<sub>3</sub>-etherate in ACN. The iodo compound is heated to 90°C for one hour with excess triethyl phosphite to produce tetraethyl 4,4'-biphenyldimethylphosphonate. Similar treatment of 1,4-naphthalene-dicarboxylic acid (Aldrich) or 9,10-anthracene-dicarboxylic acid (Aldrich) yields the respective tetraethyl phosphonates. After recrystallization from hexane, the phosphonate is



-47-

dissolved in DMF and treated with a ten-fold excess of sodium methoxide, followed by two equivalents of 5-formylsalicylic acid in DMF. After stirring at room temperature for 24 hrs, the reaction mixture is poured  
5 into water. Acidification of the water to pH 5.0 with HCl causes precipitation of the fluorescent product, 4,4'-bis(2-(3-carboxy-4-hydroxyphenyl)ethen-1-yl)-biphenyl, which can be selectively extracted into ethyl acetate from any mono-substituted by-product. Similar  
10 treatment of tetraethyl p-xylylenediphosphonate (TCI America) with 5-formylsalicylic acid, or its derivatives, gives 1,4-bis(2-(3-carboxy-4-hydroxyphenyl)ethen-1-yl)-benzene. Likewise 1,4-bis(2-(3-carboxy-4-hydroxyphenyl)ethen-1-yl)-naphthalene or 9,10-bis(2-(3-carboxy-4-hydroxyphenyl)ethen-1-yl)-anthracene is  
15 obtained by treating the appropriate phosphonate with 5-formylsalicylic acid. Other derivative are obtained by the use of other formylsalicylic acid congeners, formyl benzoic acids, or hydroxy- or methoxybenzaldehydes.

20 When backbone linkers other than those listed above are desired, the appropriate dicarboxylic acid (such as 2-bromoterephthalic acid (Aldrich)) is reduced to the diol, converted to the iodide, and then to the tetraethyl diphosphonate as described above. When side groups other  
25 than salicylic acid are desired, the appropriate phenol (which will usually also contain an acidic functionality as well) is first iodinated ortho or para (depending on the presence of other substituents) to the phenol and then formylated at the iodo-position by standard methods.

30 **Synthesis of Butadienyl (CH=CH-CH=CH) derivatives of Chrysamine-G**

p-Xylylenediphosphonic acid tetraethyl ester (TCI America, Portland, OR) is dissolved in dry DMSO and treated with a ten-fold excess of potassium t-butoxide  
35 (Aldrich), followed by two equivalents of 3-methoxycarbonyl-4-methoxycinnamaldehyde (prepared from 5-

-48-

iodo-2-methoxybenzoic acid methyl ester and acrolein by standard methods) in dry DMSO. After stirring at room temperature for 24 hrs, the reaction mixture is poured into water. Acidification of the water to pH 5.0 with HCl causes precipitation of the fluorescent product, 1,4-bis(4-(3-carboxy-4-methoxyphenyl)-1,3-butadien-1-yl)-benzene, which can be selectively extracted into ethyl acetate from any mono-substituted by-product. Cleavage of the methoxy group with excess sodium thioethoxide in refluxing DMF gives the salicylic acid derivative. Similar treatment of other combinations of backbone linker phosphonates with appropriate cinnamaldehyde derivatives as described above gives the desired products.

**Synthesis of Alkyl-Substituted Alkenyl (CR'=CR') derivatives of Chrysamine-G**

4,4'-biphenyldicarboxylic acid (Aldrich) or 1,4-benzenedicarboxylic acid (Aldrich) is first converted by reduction with  $\text{LiAlH}_4$  to the bis(hydroxymethyl) compound, which, in turn, is converted to the dicarboxaldehyde by treatment with  $\text{BaMnO}_4$  in ethyl acetate. This dialdehyde is reacted with  $\text{R}'\text{MgX}$  (where  $\text{R}'$  is a lower alkyl group and  $\text{X}$  is Br or I) via the Grignard reaction to produce  $\text{HOCR}'\text{H-Ph-Ph-CR}'\text{H-OH}$  or  $\text{HOCR}'\text{H-Ph-CR}'\text{H-OH}$ . The alkyl-substituted bis(hydroxymethyl) compound is converted to the alkyl-substituted bis(iodomethyl) compound by reaction with  $\text{NaI}$  and  $\text{BF}_3$ -etherate in ACN. The iodo compound is heated to  $90^\circ\text{C}$  for about one hour with excess triethyl phosphite to produce the alkyl-substituted tetraethyl dimethylphosphonate. Similar treatment of 1,4-naphthalene-dicarboxylic acid (Aldrich) or 9,10-anthracene-dicarboxylic acid (Aldrich) yields the respective alkyl-substituted tetraethyl dimethylphosphonates.

Alkyl-substituted alkenyl compounds of three varieties, namely  $\text{AR-CR}'=\text{CR}'-\text{Q}$ ,  $\text{AR-CR}'=\text{CH-Q}$  or  $\text{Ar-CH=CR}'-$

-49-

Q (where R' is a lower alkyl group and Q is as defined in Formula I), can then be synthesized. AR-CR'=CR'-Q compounds can be made by reacting an alkyl-substituted tetraethyl dimethylphosphonate with a suitable aryl ketone such as 5-acetylsalicylic acid (Crescent Chemical Co., Inc., Hauppauge NY). AR-CR'=CH-Q compounds can be made by reacting an alkyl-substituted tetraethyl dimethylphosphonate with a suitable aldehyde such as 5-formylsalicylic acid (Aldrich). AR-CH=CR'-Q compounds can be made by reacting tetraethyl dimethylphosphonate with a suitable aryl ketone such as 5-acetylsalicylic acid (Crescent Chemical Co., Inc., Hauppauge NY). The reaction conditions are identical to those used to make the alkenyl (CH=CH) derivatives described above.

**Synthesis of Alkynyl (C≡C) Derivatives of Chrysamine G**

5-Iodosalicylic acid (Aldrich Chemical Company, Milwaukee, WI) is converted to the methyl ester by reaction with methanol, trimethyl orthoformate and sulfuric acid. The 5-iodosalicylic acid methyl ester thus obtained is reacted with (trimethylsilyl)acetylene (Aldrich Chemical Company, Milwaukee, WI) in the presence of palladium. The trimethylsilyl group is removed and two equivalents of the resultant 5-acetylenylsalicylic acid methyl ester is reacted with 4,4'-dibromobiphenyl (Aldrich Chemical Company, Milwaukee, WI) in the presence of palladium as above. The resultant alkynyl analogue of Chrysamine G, 4,4'-bis(2-(3-methoxycarbonyl-4-hydroxyphenyl)acetylen-1-yl)-biphenyl is prepared by hydrolysis of the ester as described above.

**Alternative Synthesis of Alkynyl (C≡C) and Vinyl (CH=CH) Derivatives of Chrysamine G**

5-Bromosalicylic acid (Aldrich Chemical Company, Milwaukee, WI) is converted to the methyl ester/methyl

-50-

ether by reaction with methyl iodide in the presence of  $K_2CO_3$  as described above. The 2-methoxy-5-bromobenzoic acid methyl ester thus obtained is reacted with (trimethylsilyl)acetylene (Aldrich Chemical Company, Milwaukee, WI) in the presence of palladium. The trimethylsilyl group is removed and two equivalents of the resultant 2-methoxy-5-acetylenylbenzoic acid methyl ester is reacted with 4,4'-dibromobiphenyl (Aldrich Chemical Company, Milwaukee, WI) in the presence of palladium as above. The resultant alkynyl analogue of Chrysamine G, 4,4'-bis(2-(3-carboxy-4-methoxyphenyl)acetylen-1-yl)-biphenyl is prepared by hydrolysis of the ester as described above. This alkynyl analogue is reduced by conventional methods to form the vinyl analogue of Chrysamine G.

Another alternative synthesis of alkenyl derivatives uses either the Suzuki or Steel reactions which employs coupling of boron or tin derivatives of 2-methoxy-5-acetylenylbenzoic acid methyl ester (or other hydroxy-acid derivatives as described above) to couple to 1,4-diiodobenzene, 4,4'-diiodobiphenyl, or other di-iodo or di-bromo backbone linkers. In some cases it is preferable to make the boron or tin derivatives of the 1,4-diacetylenylbenzene (or 4,4'-diacetylenylbiphenyl) and couple these to 5-iodosalicylic acid (or similar iodo-hydroxy acids).

#### Synthesis of Di-alkynyl ( $C\equiv C-C\equiv C$ ) Derivatives of Chrysamine G

5-Iodosalicylic acid (Aldrich Chemical Company, Milwaukee, WI) is converted to the methyl ester by reaction with methanol, trimethyl orthoformate and sulfuric acid. The 5-iodosalicylic acid methyl ester thus obtained is reacted with (trimethylsilyl)acetylene (Aldrich Chemical Company, Milwaukee, WI) in the presence of palladium. The trimethylsilyl group is removed and two equivalents of the resultant 5-acetylenylsalicylic

-51-

acid methyl ester is reacted with 4,4'-bis(2-bromoacetylen-1-yl)benzene (synthesized from trimethylsilyl acetylene and 1,4-diiodobenzene) in the presence of palladium as above. The resultant di-alkynyl analogue of Chrysamine G, 4,4'-bis(4-(3-methoxycarbonyl-4-hydroxyphenyl)-1,3-butadiyn-1-yl)-benzene is prepared by hydrolysis of the ester as described above.

**Synthesis of Alkene-Alkynyl (C=C-C≡C) Derivatives of Chrysamine G**

Similar to the synthesis of alkenyl and alkynyl derivatives described above, Suzuki or Steel reactions are used to couple metal-alkenyl derivatives of benzene, biphenyl, other other backbone groups to 2-methoxy-5-(2-iodoacetylen-1-yl)-benzoic acid methyl ester (other other appropriate hydroxy acid derivatives) or alternatively, 5-metal-alkenyl derivatives of 2-methoxybenzoic acid methyl ester can be coupled to 1,4-bis(2-iodoacetylen-1-yl)-benzene (or 4,4'-bis(2-iodoacetylen-1-yl)-biphenyl).

**Synthesis of Alkyl (CH<sub>2</sub>-CH<sub>2</sub>) or Di-Alkyl (CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>) Derivatives of Chrysamine G**

Either the alkenyl, alkynyl, di-alkenyl, di-alkynyl or alkene-alkynyl derivatives described above are hydrogenated by standard methods using hydrogen gas and a platinum or palladium catalyst.

-52-

**Synthesis of Di-fluoro Alkenyl Chrysamine  
G Derivative**

The 5-fluoro derivative, 1,4-bis(2-(2-hydroxy-3-carboxy-5-fluorophenyl)ethen-1-yl)-benzene), is  
5 synthesized by substituting 3-formyl-5-fluorosalicylic acid for 5-formyl-salicylic acid. [ $^{18}\text{F}$ ]aryl fluorides derivatives of Chrysamine G can be prepared by substituting  $^{18}\text{F}$ -labeled precursors such as [ $^{18}\text{F}$ ]LiBF<sub>4</sub>, in the Schiemann reaction, via triazene decomposition with  
10 Cs [ $^{18}\text{F}$ ], or via nucleophilic  $^{18}\text{F}$ -for-X substitution, where X = tosyl, triflate, NO<sub>2</sub>, <sup>+</sup>N(CH<sub>3</sub>)<sub>3</sub>, or halogen. See Fowler, J. and Wolf, A. in POSITRON EMISSION TOMOGRAPHY AND AUTORADIOGRAPHY (Phelps, M., Mazziota, J., and Schelbert, H. eds.) 391-450 (Raven Press, NY, 1986) and  
15 Kilbourn, M. Fluorine-18 labeling of radiopharmaceuticals. (Natl. Acad. Press, Washington, D.C.) (1990).

**Synthesis of Aromatic Fluoroalkyl and Fluoroalkoxy Derivatives**

20 Aromatic fluoroalkyl derivatives are synthesized employing the method of Bishop et al., *J. Med. Chem.* 34: 1612 (1991) in which Claisen rearrangement of the appropriate O-allyl ethers forms an aromatic allyl derivative which can be further functionalized to yield  
25 the fluoroethyl or fluoropropyl derivatives. Alternatively, an aromatic iodide can be readily converted to an aromatic alkyne consisting of two to five carbon atoms in length using the palladium-assisted coupling methodology of Sonogashira et al., *Tetrahedron Letters* 4467-4470 (1975). Subsequent derivatization of  
30

-53-

the alkyne yields the fluoroalkyl derivative.

Fluoroalkoxy derivatives may be prepared by the method of Chumpradit et al., *J. Med. Chem.* 36: 21 (1993) in which alkylation of the appropriate phenol with the appropriate 1-bromo (or iodo or sulfonyloxy)-omega-fluoroalkane yields the corresponding fluoroalkoxy derivative.

#### **Radiofluorination of Aromatic Alkylsulfonyloxy and Alkoxy sulfonyloxy Derivatives**

Radiofluorination to yield the aromatic [ $^{18}\text{F}$ ]fluoroalkyl and [ $^{18}\text{F}$ ]fluoroalkoxy derivatives is performed by the method of Mathis et al., *Nucl. Med. Biol.* 19: 571 (1992) in which aromatic alkyl- or alkoxy sulfonyloxy (e.g. alkoxytosylate) derivatives are substituted with [ $^{18}\text{F}$ ]fluoride to yield aromatic [ $^{18}\text{F}$ ]fluoroalkyl and [ $^{18}\text{F}$ ]fluoroalkoxy compounds.

#### **Radio-Iodination and Radio-Bromination by the Tri-Alkyl Tin Route**

##### Synthesis of Tri-Alkyl Tin Derivatives

The general structure of the alkenyl tri-alkyl tin derivative of Chrysamine G is shown in Figure 2B. In general, one tri-alkyl tin group will be substituted at the 3-position on one side of the biphenyl moiety, but other positions, including the salicylic acid or heterocyclic moiety are also potential targets. These tri-alkyl tin derivatives are stable immediate precursors for preparation of the radioiodinated and radiobrominated compounds to be used in humans. More specifically, these tri-alkyl tin derivatives are used to prepare the halogenated radioactive compounds applicable for use in *in vivo* imaging of amyloid.

-54-

**General Procedures for the Synthesis of Tri-Alkyl Tin Derivatives**

Tri-alkyl tin derivatives are prepared from the appropriate arylhalides,  $[(C_6H_5)_3P]_3Pd(0)$ , and hexaalkylditin by previously published procedures including Kosugi, M., et al., *Chem. Lett.* 1981: 829; Heck, R. *Pure and Appl. Chem.* 1978: 691; Echavarren, A. and Stille, J. *J. Am. Chem. Soc.* 1987: 5478; Mitchell, T. *J. Organometallic Chem.* 1986: 1; and Stille, J. *Pure and Applied Chem.* 1985: 1771. These derivatives also can be obtained by the use of *n*-BuLi and trialkyl tin chloride by the procedure of Mathis et al., *J. Labell. Comp. and Radiopharm.* 1994: 905.

**Synthesis of the 3-trialkyl tin derivative of 4,4'-bis(3-methoxycarbonyl-4-hydroxyphenylazo)-biphenyl**

3-Bromo or 3-iodo-4,4'-bis(3-methoxycarbonyl-4-hydroxyphenylazo)-biphenyl or its dimethyl ether are prepared by synthesis of 3-bromo- or 3-iodobenzidine (see above), tetra-azotization and coupling to methyl salicylate as for the synthesis of Chrysamine G, and methylation of the phenol as described above when the methoxy compound is desired. Under an argon atmosphere, 1 mmol of the phenolic ester or the methoxy ester,  $[(C_6H_5)_3P]_3Pd(0)$  (0.1 to 0.2 mmol), hexabutylditin or hexamethyl ditin (1.25 mmol), and dioxane (25 ml) is heated at 70°C for 16 hrs. The reaction mixture is cooled and the solvent is evaporated. Tri-alkyl tin halide is removed with aqueous KF. The organics are extracted with ethyl acetate, dried over magnesium sulfate, filtered, and the solvent is evaporated under reduced pressure. The residue is purified on silica gel to obtain 3-trialkyltin-4,4'-bis(3-methoxycarbonyl-4-hydroxyphenylazo)-biphenyl.

**Radio-Iodination or Radio-Bromination of Tri-Alkyl Tin Derivatives**

The tributyl or trimethyl tin derivatives are radio-



-55-

iodinated with Na[<sup>125</sup>I] or Na[<sup>123</sup>I] or radio-brominated with Na[<sup>75</sup>Br] or Na[<sup>76</sup>Br] by published procedures such as Mathis et al., *J. Labell. Comp. and Radiopharm.* 1994: 905; Chumpradit et al., *J. Med. Chem.* 34: 877 (1991);  
5 Zhuang et al., *J. Med. Chem.* 37: 1406 (1994); Chumpradit et al., *J. Med. Chem.* 37: 4245 (1994). In general, 0.5 mg of tri-alkyl tin compound, 0.2 ml of anhydrous acetonitrile, 10 µl of 2M H<sub>3</sub>PO<sub>4</sub>, 2-100 µl of a solution of high specific activity (>2000 Ci/mmol) Na[<sup>125</sup>I] or Na[<sup>123</sup>I]  
10 (or Na[<sup>75</sup>Br] or Na[<sup>76</sup>Br]) in pH 9-12 NaOH, and dichloramine-T (DCT) (20 µl of 2.5 mg/ml DCT in acetonitrile) are placed in a 1 ml Reacti-Vial. The vial is capped and the mixture is stirred at room temperature in the dark. The reaction is monitored by HPLC and after  
15 30 min is quenched with 50 µl of 2 M Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>. The product is purified by standard chromatographic techniques. Mathis et al., *J. Labell. Comp. and Radiopharm.* 1994: 905. Similarly, low specific activity <sup>18</sup>F derivatives are prepared by analogous procedures.

20       **General Procedures for the Preparation of Non-Radioactive I, Br, Cl, F and -SH Derivatives**

In general, 3- or 4-amino derivatives of 5-formylsalicylic acid, or the corresponding derivatives of the heterocyclic analogues of salicylic acid shown in  
25 Figure 2, are converted to the corresponding diazo compounds with sodium nitrite and HCl or H<sub>2</sub>SO<sub>4</sub>. The iodine derivatives are directly prepared by forming the diazonium iodide which is then converted into the aryl iodide, or by way of the triazene intermediates. See,  
30 e.g., Greenbaum, F. *Am. J. Pharm.* 108: 17 (1936), Satyamurthy, N. and Barrio, J., *J. Org. Chem.* 48: 4394 (1983) and Goodman, M. et al., *J. Org. Chem.* 49: 2322 (1984). Aryl bromides and chlorides are prepared from the diazo compounds by treatment with CuCl or CuBr  
35 according to the Sandmeyer reaction or via the triazene as for the iodine derivatives. Aryl fluorides are

-56-

prepared by treating the diazonium compounds with  $\text{NaBF}_4$ ,  $\text{HBF}_4$ , or  $\text{NH}_4\text{BF}_4$  according to the Schiemann reaction or via triazene decomposition similar to the iodine derivatives. Aryl thiols are prepared from the diazonium compounds by treatment with sulfur-containing nucleophiles such as  $\text{HS}^-$ ,  $\text{EtO-CSS}^-$ , and  $\text{S}_2^{2-}$ . Alternatively, aryl thiols can be prepared by replacement of aryl halides with sulfur containing nucleophiles. These reactions are described in March, J., *ADVANCED ORGANIC CHEMISTRY: REACTIONS, MECHANISMS, AND STRUCTURE* (3rd Edition, 1985).

#### General Procedures for the Preparation of Radioactive C, F and Tc Derivatives

In addition to the above procedures, high specific activity radiolabeling with  $^{99\text{m}}\text{Tc}$  for SPECT or with the positron-emitting radionuclides  $^{11}\text{C}$ ,  $^{18}\text{F}$ ,  $^{75}\text{Br}$  and  $^{76}\text{Br}$  is accomplished according to literature-based methods well known in the art. Some of the potential specific methods are described below, but there are other well-known methods which will be apparent to those skilled in the art and are described in Fowler, J. and Wolf, A. *Positron emitter-labeled compounds* in *POSITRON EMISSION TOMOGRAPHY AND AUTORADIOGRAPHY* (Phelps, M., Mazziota, J., and Schelbert, H. eds.) p 391-450 (Raven Press, NY) (1986), Coenen, H. et al., *Radiochimica Acta* 34: 47 (1983), and Kulkarni, *Int. J. Rad. Appl. & Inst. (Part B)* 18: 647 (1991), the contents of which are hereby incorporated by reference.

$^{99\text{m}}\text{Tc}$  derivatives are prepared by complexation with the aryl thiols. Radiolabeling with  $^{11}\text{C}$  can be readily done via N-methylation, O-methylation as described above substituting  $[^{11}\text{C}]$ methyl iodide,  $[^{11}\text{C}]$ alkylation, or  $[^{11}\text{C}]$ carboxylation of suitable alkyl, alkenyl, or alkynyl Chrysamine G analogues.  $[^{18}\text{F}]$ aryl fluorides derivatives can be prepared by substituting  $^{18}\text{F}$ -labeled precursors such as  $[^{18}\text{F}]\text{LiB}_4$  in the Schiemann reaction described above, via triazene decomposition with  $\text{Cs}[^{18}\text{F}]$ , or via

-57-

nucleophilic  $^{18}\text{F}$ -for-X substitution, where X = tosyl, triflate,  $\text{NO}_2$ ,  $^+\text{N}(\text{CH}_3)_3$ , or halogen. Radiobromination using  $^{75}\text{Br}$  and  $^{76}\text{Br}$  can be accomplished using either electrophilic ( $\text{Br}^+$ ) or nucleophilic ( $\text{Br}^-$ ) substitution techniques analogue to radioiodination techniques, see Coenen, H., *supra*.

**Synthesis of the 3-Hydroxy-1,2-benzisoxazole derivative and related derivatives (see Figure 2C)**

2,6-Dihydroxybenzoic acid ( $\gamma$ -resorcylic acid) methyl ester (TCI America, Portland, OR) is converted to the hydroxamic acid by the use of hydroxylamine hydrochloride according to the method of Böshagen (Chem. Ber 100: 954-960; 1967). The hydroxamic acid is converted to the corresponding 3-hydroxy-1,2-benzisoxazole with the use of  $\text{SOCl}_2$  and then triethylamine, also by the method of Böshagen (Chem. Ber 100: 954-960; 1967). This compound is then converted to the formyl derivative and coupled to the appropriate tetraethyl diphosphonate (which in some cases may be brominated) as described above under the synthesis of alkenyl derivatives. The bromo derivatives can then be converted to the tri-alkyl tin and iodo-derivatives as described above.

Alternatively, 5-formyl salicylic acid is coupled to the appropriate tetraethyl diphosphonate as usual and the resulting 4,4'-bis(2-(3-carboxy-4-hydroxyphenyl)ethen-1-yl)-benzene is converted first to the dimethyl ester and then to the hydroxamic acid and finally the benzisoxazole by the method of Böshagen as described above. A third type of 3-hydroxy-1,2-benzisoxazole is synthesized from several isomeric dihydroxy benzenedicarboxylic acids including 4,6-dihydroxy-1,3-benzenedicarboxylic acid, 3,6-dihydroxyphthalic acid, and 2,5-dihydroxyterephthalic acid (Aldrich Chem. Co., Milwaukee, WI). After formylation and coupling to the appropriate tetraethyl diphosphonate by standard procedures, followed

-58-

by conversion to the dimethyl esters, the dihydroxy/diesters are converted to dihydroxy/dihydroxamic acids by reaction with hydroxylamine by the method of Böshagen described above.

- 5 Conversion to the double benzisoxazole is effected by treatment with  $\text{SOCl}_2$  and triethylamine, again, by the method of Böshagen described above.

**Synthesis of the phthalimide or isoindole-1,3(2H)-dione derivative (see Figure 2D)**

- 10 3-Hydroxyphthalimide made from 3-hydroxyphthalic anhydride (Aldrich Chemical Company, Milwaukee, WI) is converted to the formyl derivative and coupled to the appropriate tetraethyl diphosphonate (which in some cases may be brominated) as described above under the synthesis  
15 of alkenyl derivatives. The bromo derivatives can then be converted to the tri-alkyl tin and iodo-derivatives as described above.

**Synthesis of the phthalhydrazide or 2,3-benzodiazine-1,4(2H,3H)-dione derivative (see Figure 2E)**

- 20 3-Hydroxyphthalhydrazide made from the reaction of 3-hydroxyphthalic anhydride (Aldrich Chemical Company, Milwaukee, WI) with hydrazine is converted to the formyl derivative and coupled to the appropriate tetraethyl  
25 diphosphonate (which in some cases may be brominated) as described above under the synthesis of alkenyl derivatives. The bromo derivatives can then be converted to the tri-alkyl tin and iodo-derivatives as described above.

- 30 **Synthesis of the 2,3-benzoxazine-1,4(3H)-dione derivative (see Figure 2F)**

- 3-Hydroxyphthalic anhydride (Aldrich Chemical Company, Milwaukee, WI) is converted to the 2,3-benzoxazine with the use of hydroxylamine. The  
35 benzoxazine derivative is then converted to the formyl

-59-

derivative and coupled to the appropriate tetraethyl diphosphonate (which in some cases may be brominated) as described above under the synthesis of alkenyl derivatives. The bromo derivatives can then be converted to the tri-alkyl tin and iodo-derivatives as described above.

**Synthesis of the (2H)1,3-benzoxazine-2,4(3H)-dione derivative (see Figure 2G)**

This compound is synthesized by the method of Effenberger et al., (Chem. Ber. 105: 1926-1942; 1972). Briefly, 4-hydroxybenzaldehyde (Aldrich) is coupled with the appropriate tetraethyl diphosphonate via the same procedure used for formylsalicylic acid derivatives. This adduct is then converted to the carbamate by reaction with ethoxycarbonylisocyanate ( $O=C=N-CO-O-Et$ ) in the presence of triethylamine. This substituted carbamate (or N-ethoxycarbonyl-carbamic acid-phenyl ester) is converted to the benzoxazinedione by heating in diphenyl ether. The benzoxazinedione is then converted to the tri-alkyl tin and iodo-derivatives, as described above.

**Synthesis of the (3H)2-benzazine-1,3(2H)-dione derivative (see Figure 2H)**

3-Hydroxyphenylacetic acid (Aldrich Chemical Company, Milwaukee, WI) is formylated, converted to the amide and then coupled with the appropriate tetraethyl diphosphonate via the same procedure used for formylsalicylic acid derivatives. This adduct is then converted to the N-(3-hydroxyphenylacetoxyl)-carbamic acid ethyl ester derivative by reaction with ethyl chloroformate. This substituted carbamate is converted to the benzazinedione by heating in diphenyl ether. The benzazinedione is then converted to the tri-alkyl tin and iodo-derivatives, as described above.

-60-

**Synthesis of the 1,8-Naphthalimide derivative  
(See Figure 2I)**

4-Amino-1,8-naphthalimide is converted to the formyl derivative and coupled to the appropriate tetraethyl  
5 diphosphonate (which in some cases may be brominated) as described above under the synthesis of alkenyl derivatives. The bromo derivatives can then be converted to the tri-alkyl tin and iodo-derivatives as described above.

10 **Synthesis of Tetrazole and Oxadiazole Derivatives  
(See Figure 2J and 2K)**

2-Cyanophenol (Aldrich Chemical Company, Milwaukee, WI) is converted to the tetrazole by reaction with sodium or aluminum azide according to the method of Holland and  
15 Pereira *J. Med. Chem.* 10: 149 (1967) and Holland U.S. Patent No. 3,448,107. Briefly, 2-cyanophenol or alkenyl cyanophenol derivatives of Chrysamine G (made by coupling 4-formyl-2-cyanophenol with the appropriate tetraethyl diphosphonate) (1 mmol) in 40 ml DMF is treated with  
20 sodium azide (10 mmol) and triethylamine hydrochloride (10 mmol) under argon. The mixture is stirred at 120°C for 2 hrs after which the mixture is cooled and worked up in a manner analogous to that described above for Chrysamine G.

25 The oxadiazoles are synthesized by treatment of the tetrazoles prepared as above with an acid anhydride (such as acetic anhydride). An alternate method is that of Bamford et al., *J. Med. Chem.* 38: 3502 (1995). In this procedure, hydrazide alkenyl derivatives of Chrysamine G  
30 or salicylic acid (obtained by treatment of the respective esters with hydrazine) are treated with methyl isothiocyanate in the presence of dicyclohexylcarbodiimide.

-61-

### Synthesis of Chrysamine G Derivatives for Use as Controls

The aniline derivative is synthesized by substituting two equivalents of aniline (Fisher Chemical Co., Fair Lawn, NJ) for each equivalent of benzidine. 5 The 2,2'-disulfonic acid derivative is synthesized by substituting benzidine-2,2'-disulfonic acid (Pfaltz & Bauer, Inc., Waterbury, CT) for benzidine. The phenol derivative is synthesized by substituting one equivalent 10 of phenol for each equivalent of salicylic acid. Congo red (Aldrich certified grade) is obtained commercially.

### EXAMPLE 2. CHRYSAMINE G AND CHRYSAMINE G DERIVATIVES BIND SPECIFICALLY TO A $\beta$

#### Binding to synthetic A $\beta$ (10-43)

15 Chrysamine G binds well to synthetic A $\beta$ (10-43) peptide *in vitro*. Figure 4A shows a Scatchard analysis of the binding of Chrysamine G to A $\beta$ (10-43). The higher affinity component has a  $K_D$  of 0.257  $\mu$ M and a  $B_{max}$  of 3.18 nmoles Chrysamine G/mg A $\beta$ (10-43). The lower affinity 20 component is less well defined by these data, but appears to have a  $K_D$  of 4.01  $\mu$ M and a  $B_{max}$  of 18.7 nmoles Chrysamine G/mg A $\beta$ (10-43). The low affinity component represents the binding of Chrysamine G at high concentrations to a distinct, low-affinity site, not the 25 binding to an impurity in the preparation. The amount of Chrysamine G injected *in vivo* is so low that there is not any binding to the low-affinity component. At very low concentrations, the ratio of high-to-low affinity binding is very large.

30 The amount of Chrysamine G binding is linear with peptide concentration over the range employed, as shown in Figure 5.

-62-

### Kinetics of Binding

Kinetic studies showed a fairly rapid association (Figure 6A), essentially complete by 1 min, at a [Chrysamine G] = 112  $\mu$ M with a  $t_{1/2}$  of  $8.9 \pm 1.8$  sec and a somewhat less rapid dissociation (Figure 6C),  $t_{1/2} = 55 \pm 9.4$  sec [dissociation rate constant ( $k_{-1}$ ) =  $1.26 \times 10^{-2}$  sec $^{-1}$ ]. Figure 6B shows a transformation of the association kinetic data according to the method of Bennett and Yamamura. Bennett, J.P. and Yamamura, H.I. in NEUROTRANSMITTER RECEPTOR BINDING (N.Y.: Raven Press 1985) pp. 61-89. The linear portion of the association curve in Figure 6A is transformed into the line of Figure 6B, in which  $\ln[B_{eq}/(B_{eq}-B_t)]$  is plotted versus time, where  $B_{eq}$  is the amount of Chrysamine G bound at equilibrium (4 min) and  $B_t$  is the amount bound at time=t. The slope of this line equals  $k_{observed}$  and  $k_1 = (k_{observed} - k_{-1})/[Chrysamine G]$ , where  $k_{-1}$  is the dissociation rate constant determined from the data in Figure 6C. The curve in Figure 6C follows the equation:

$$A_t = A_0 e^{-k_{-1}t} \quad \text{where}$$

$A_t$  is the amount of Chrysamine G remaining bound at time=t,  $A_0$  is the amount of Chrysamine G bound at time=0,  $t$  is the time in min, and  $k_{-1}$  is the dissociation rate constant. From this analysis, the association rate constant ( $k_1$ ) is calculated to be  $3.75 \times 10^4$  M $^{-1}$  sec $^{-1}$  giving a  $K_D = k_{-1}/k_1 = 0.34$   $\mu$ M, in good agreement with the Scatchard analysis.

### Chrysamine G Derivatives Can Inhibit the Binding of Chrysamine G to A $\beta$

$K_i$  values for the inhibition of [ $^{14}$ C]Chrysamine G binding to A $\beta$ (10-43) by the Chrysamine G analogues are shown under the chemical structures in Figure 1 and several displacement curves are shown in Figure 3.  $K_i$  is defined as  $IC_{50}/(1+[L]/K_D)$ , where [L] is the concentration



-63-

of [ $^{14}\text{C}$ ]Chrysamine G in the assay (.100-.125  $\mu\text{M}$ ) and  $K_D$  is 0.26  $\mu\text{M}$ , the  $K_D$  of Chrysamine G determined by the Scatchard analysis above. Chrysamine G itself gives a  $K_1$  of  $0.37 \pm 0.04$   $\mu\text{M}$ , a value very consistent with those  
5 obtained from the Scatchard and kinetic analyses. Congo red gives a  $K_1$  of  $2.82 \pm .84$   $\mu\text{M}$ . The difluoro derivative of Chrysamine G, (5-FSA)CG, (Figure 1) is one-third as potent as Chrysamine G itself ( $K_1 = 1.16 \pm 0.19$   $\mu\text{M}$ ). The activity of the difluoro Chrysamine G derivative suggests  
10 that an  $^{18}\text{F}$  difluoro Chrysamine G derivative works for PET imaging and an  $^{19}\text{F}$  difluoro Chrysamine G derivative works for MRS/MRI imaging of brain.

The 3-ICG is slightly more potent than Chrysamine G. The activity of the 3-ICG derivative suggests that an  $^{123}\text{I}$   
15 difluoro Chrysamine G derivative works for SPECT imaging. Methylating the phenol of 3-ICG decreases the affinity by a factor of 10 in 3-ICG(OMe) $_2$ . Methylating the carboxylate group effected an even greater (about 200-fold) decrease in affinity in CG(COOMe) $_2$ . Removing the  
20 acid moiety entirely, as in the phenol derivative, completely destroyed binding affinity.

These results suggest that the acid moiety of Chrysamine G analogues plays the major role in binding to A $\beta$  and that the phenol moiety plays a facilitating role.  
25 The effect of the phenol could occur through hydrogen bonding to the acid which could serve to stabilize the structural orientation of the acid moiety. The presence of a phenol in the ortho position could also alter the charge distribution of the acid either through hydrogen  
30 bonding or through changes in the charge distribution of the aromatic system as a whole. Alternatively, the phenol could directly participate in binding to the amyloid via a bi-dentate attachment of both the phenol and the acid to the amyloid binding site. Adding a  
35 second phenol ortho to the carboxylate as in the resorcylic acid derivative, (6-OHSA)CG, produces the highest affinity compound in this series having a  $K_1$  of

-64-

0.094 ± .02 μM.

Increasing the lipophilicity of the biphenyl backbone appears to increase the affinity somewhat. The di-halo derivatives, 3,3'-I<sub>2</sub>CG, 3,3'-Br<sub>2</sub>CG, and 3,3'-Cl<sub>2</sub>CG, all have very similar K<sub>i</sub> values which are about half that of Chrysamine G.

Distorting the dihedral angle between the phenyl rings of the biphenyl group by substitution at the 2-position markedly diminishes affinity. This is demonstrated by the inactivity of the 2,2'-di-sulfonic acid derivative of Chrysamine G, 2,2'-(SO<sub>3</sub>)<sub>2</sub>CG. Since the 3,3' di-carboxylic derivative, 3,3'-(COOH)<sub>2</sub>CG, shows only a 7-fold loss of activity from Chrysamine G, it is unlikely that the additional acidic moieties are the sole cause for the loss of activity in the 2,2'-disulfonic acid. This 2,2'- derivative is unique in that the bulky sulfonate groups in the 2-position force the biphenyl group out of planarity. Molecular modelling studies showed that the dihedral angle between the two biphenyl benzene rings in the 2,2'-disulfonic acid derivative is 83°. This angle is approximately 35-40° in Chrysamine G and all of the other active derivatives.

In an attempt to explore the importance of the bidentate nature of the functional groups of Chrysamine G, the binding of an aniline derivative which represents one-half of a Chrysamine G molecule (Figure 1) was studied. An approximation of the energy of binding can be calculated from the equation:

$$\Delta G = -RT \ln K_{eq}$$

wh

ere ΔG is the energy of the binding reaction, R is the molar gas constant [8.31441 J/(mole·°K)], T is temperature in °K and K<sub>eq</sub> is the equilibrium constant for the reaction:



and K<sub>eq</sub> = 1/K<sub>D</sub> ≈ 1/K<sub>i</sub>. Using the value of 0.26 μM for the K<sub>D</sub>

-65-

of Chrysamine G, the energy of binding is roughly 38 KJ/mole. If the aniline derivative binds with one-half of this energy, the expected energy of binding would be about 19 KJ/mole. From the  $K_1$  of 73  $\mu$ M for the aniline derivative, the energy of binding is 23 KJ/mole which is in acceptable agreement with the predicted value. The importance of the hydrophobic region of Chrysamine G and the aniline derivative is demonstrated by the total lack of binding activity of salicylic acid itself.

10       The affinity of Chrysamine G for A $\beta$  appears to be several fold greater than the affinity of Congo red for this peptide. The binding is reversible with a dissociation constant of approximately 250-400 nM, whether measured by Scatchard analysis, kinetic methods,  
15       or inhibition of binding. Owing to the non-crystalline, poorly soluble nature of amyloid fibrils, the structure of Congo red or Chrysamine G complexes with amyloid has never been defined by precise structural techniques such as x-ray crystallography or multi-dimensional NMR.  
20       Models of Congo red interactions with amyloid have been proposed. Cooper, *Lab. Invest.* 31: 232 (1974); Romhanyi, *Virchows Arch.* 354: 209 (1971). This work suggests that Congo red does not bind to a single amyloid peptide molecule, but binds across several A $\beta$  molecules oriented  
25       by virtue of the beta-sheet fibril. Klunk et al., *J. Histochem. Cytochem.* 37: 1273 (1989).

      Figure 7 shows a schematic of this model, generated using MacroModel 2.5, in which Chrysamine G spans 5 peptide chains in an anti-parallel beta-sheet  
30       conformation. The peptides are used without further structural refinement. The peptides are aligned so that alternate chains were spaced 4.76 Å apart, characteristic of beta-sheet fibrils. Alternate peptide chains are drawn in black and white. Chrysamine G (black) is energy  
35       minimized and aligned with the fibril model to maximize contact with lysine-16 (light grey ovals in top figure) and the hydrophobic phenylalanine 19/20 region (bottom).

-66-

The two views are of the same model at approximately 90° from one another. The white arrows indicate the direction taken to obtain the alternate view.

5 The 19.1 Å spacing between the carboxylic acid moieties of Chrysamine G matches well with the distance of 19.0 Å across the 5 chains (4 x 4.76 Å between adjacent chains shown by Kirschner et al., *Proc. Natl. Acad. Sci. U.S.A.* 83: 503 (1986)). If the native structure of Aβ involves a hairpin loop structure as  
10 Hilbich et al., suggest (Hilbich et al., *J. Mol. Biol.* 218: 149 (1991)), then chains 1 and 2, 3 and 4, 5 and 6, etc., would be folded halves of the same molecule, but the model would otherwise be the same. Also important to note is the necessity for positively charged amino acid  
15 residues in this model, such as lysine-16 in Aβ. Previous work has shown that Congo red binding correlates well with the number of positively charged amino acids in a sample of amyloid fibrils. Klunk et al., *J. Histochem. Cytochem.* 37: 1273 (1989). The bidentate nature of the  
20 model in Figure 7 and the importance of hydrophobic interactions is supported by the decrease in affinity of the mono-dentate aniline analogue of Chrysamine G and the inactivity of salicylic acid as well as the increased potency of the more lipophilic compounds having two  
25 halogens on the benzidine moiety (see Figure 1). The importance of the nearly planar biphenyl group is suggested by the inactivity of the 2,2'-disulfonic acid derivative.

30 **EXAMPLE 3.      CHRYSAMINE G DISTINGUISHES ALZHEIMER'S DISEASE BRAIN FROM NORMAL BRAIN**

**Characterization of the Binding of  
Chrysamine G to AD Brain**

Scatchard analyses of the binding of Chrysamine G and Chrysamine G derivatives to AD brain samples were  
35 performed in an effort to understand the increased binding of Chrysamine G to AD brain. (Figure 4B, Table

-67-

1). Under the conditions employed, control and AD brain showed a single binding component. The  $K_D$  in AD brain was 16% lower than control but the difference was not significant ( $p=.29$ ). The  $B_{max}$  in AD brain was 36% higher than the  $B_{max}$  in control brain, but, again, the difference did not reach significance ( $p=.09$ ). Therefore, the increased binding in AD brain appears to be mainly due to the presence of more of the same binding component which exists in control brain, rather than the presence of a unique component.

Table 1. Comparison of binding Parameters in AD and control brain

|                  | $K_D$ ( $\mu M$ ) | $B_{max}$<br>( $pmol/\mu g$<br>prot) |
|------------------|-------------------|--------------------------------------|
| Control<br>(n=6) | $0.47 \pm .049$   | $0.576 \pm .092$                     |
| AD (n=5)         | $0.39 \pm .048$   | $0.784 \pm .061$                     |

The binding of CG to AD brain significantly correlated with numbers of NPs in the association cortices of the brain. Figure 8A shows the correlation of [ $^{14}C$ ]CG binding with numbers of NPs in the superior/middle frontal and superior temporal cortex of AD brain. The correlation with NPs was significant whether controls were included ( $r=0.69$ ;  $p=0.001$ ) or if the AD brains were considered alone ( $r=0.59$ ;  $p=0.007$ ). Figure 8B shows a similar correlation with NFT counts. As with NPs, the correlation with NFTs was significant whether controls were included ( $r=0.60$ ;  $p=0.001$ ) or if the AD brains were considered alone ( $r=0.50$ ;  $p=0.026$ ). The correlation with NFTs is not surprising since CG is a derivative of Congo red, which stains NFTs. The number of NPs was significantly correlated with the number of NFTs ( $r=0.82$ ;  $p=.0001$ ).

-68-

Only qualitative data on the presence or absence of amyloid angiopathy was available for the brains used in this study, so similar correlations could not be performed between CG binding and cerebrovascular amyloid levels. The presence of amyloid angiopathy does appear to be a confounding variable in the correlation of CG binding with NP counts. Figure 8A shows the improved correlation of CG binding to NP counts in brains without amyloid angiopathy ( $r=0.79$ ;  $p=0.01$ ) compared to those brains with cerebrovascular amyloid deposits ( $r=0.49$ ;  $p=0.15$ ). A similar improvement was not found in the correlation to NFT counts.

The  $K_D$  for [ $^{14}\text{C}$ ] Chrysamine G binding to AD brain is similar to that found for [ $^{14}\text{C}$ ]Chrysamine G binding to synthetic A $\beta$  *in vitro*, suggesting that binding in brain homogenates also may represent interaction with A $\beta$ . The correlation of Chrysamine G binding to NFTs may indicate that Chrysamine G binds to these structures in brain homogenates as well. Alternatively, since the number of NFTs correlates closely with the number of NPs, the correlation of [ $^{14}\text{C}$ ]Chrysamine G binding to NFTs may just be an epiphenomenon of Chrysamine G binding to NPs.

The useful Chrysamine G derivatives or analogues provided herein have binding affinities that are at least in the range of 0.01 to 10.0  $\mu\text{M}$   $K_D$ , as measured by binding to either synthetic A $\beta$  peptide or Alzheimer's Disease brain tissue; higher affinity compounds having binding affinities in the range of 0.0001 to 0.01  $\mu\text{M}$  are also useful in the method of the present invention.

Considering the above, Chrysamine G binding may not be specific for A $\beta$ . Instead, Chrysamine G binding may reflect the total amyloid "load" of the brain, comprised of aggregated deposits of A $\beta$  in neuritic plaques and cerebrovascular amyloid. Deposits of

-69-

phosphorylated tau protein in NFTs may contribute to Chrysamine G binding as well. Goedert, M. et al., *PNAS* 85: 4051 (1988). NFTs also are composed of anti-parallel beta-sheet fibrils similar in quaternary structure to fibrils of A $\beta$ . Kirschner et al., *Proc. Natl. Acad. Sci. U.S.A.* 83: 503 (1986).

#### **Total and Relative Chrysamine G Binding Distinguishes AD From Normal Brain**

*In vitro* binding assays such as those described above and below are widely used in the art as models to screen compounds for *in vivo* binding in brain and to predict success in subsequent *in vivo* imaging studies. See, Young, A. et al., *Receptor Assays: In Vitro and In Vivo*. in *POSITRON EMISSION TOMOGRAPHY AND AUTORADIOGRAPHY* (Phelps, M., Mazziota, J., and Schelbert, H. eds.) pp. 73-111 (1986). The labeled Chrysamine G and Chrysamine G derivatives of the invention also may be used in the *in vitro* binding assays described above and below to quantitate amyloid deposits in biopsy or post-mortem specimens.

Saturable (specific) binding of [ $^{14}\text{C}$ ] Chrysamine G was observed both in AD brain and control brain homogenates and constituted 60-80% of total binding in AD brain. Non-saturable binding was very similar in AD and control brain. Both saturable and total binding were greater in AD brain than in control. Despite the lower sensitivity obtained when using total binding, this parameter is more predictive of success in *in vivo* studies which are the ultimate goal of this invention. Also for the purpose of extension to *in vivo* studies, it is advantageous if Chrysamine G binding in cortical areas is normalized to a brain area in which Chrysamine G binding is very similar in both AD and control brain. This obviates the need to calculate the absolute quantity bound which is difficult to do *in vivo*. We examined binding in the cerebellum as a potential

control area because classical NPs are exceedingly rare in this brain area (Joachim et al., *Am. J. Pathol.* 135: 309 (1989)).

The average amount of [<sup>14</sup>C]Chrysamine G bound to control cerebellum is nearly identical to the amount bound to AD cerebellum (Table 2), supporting the use of cerebellum as an internal control. Therefore, the cerebellar ratio (CBR) accurately reflects the absolute quantity of [<sup>14</sup>C]Chrysamine G bound and offers the advantage of providing an internal control for each brain. Binding is greater in AD brain whether expressed in absolute terms of fmol/ $\mu$ g protein (Table 2) or as a ratio to the binding in the cerebellum of the same brain (Table 3). The CBR is the more sensitive measure and shows less variability between brains. The use of total binding and CBRs greatly facilitates extension of these *ex vivo* results to *in vivo* studies. Accordingly, the results below are expressed in these terms whenever appropriate.

**Table 2. Comparison of total binding in AD and control brain\*.**

| Brain Area              | Control<br>(CBR)     | AD<br>(CBR)           | p Value |
|-------------------------|----------------------|-----------------------|---------|
| Frontal Pole            | 0.87 $\pm$ .04 (n=6) | 1.87 $\pm$ .25 (n=10) | p<0.004 |
| Superior/Middle Frontal | 0.73 $\pm$ .02 (n=8) | 1.84 $\pm$ .18 (n=11) | p<0.001 |
| Superior Temporal       | 0.86 $\pm$ .08 (n=8) | 1.63 $\pm$ .17 (n=11) | p<0.002 |
| Head of Caudate         | 0.95 $\pm$ .04 (n=4) | 1.76 $\pm$ .31 (n=7)  | p<0.04  |
| Inferior Parietal       | 0.90 $\pm$ .08 (n=8) | 1.93 $\pm$ .20 (n=11) | p<0.001 |
| Occipital               | 0.77 $\pm$ .13 (n=8) | 1.44 $\pm$ .20 (n=11) | p<0.02  |

\*High- and low-plaque AD brains combined.



Table 3. Comparison of total binding in AD and control brain as a ratio to cerebellum\*.

| Brain Area              | Control<br>(fmol/ $\mu$ g protein) | AD<br>(fmol/ $\mu$ g protein) | p Value |
|-------------------------|------------------------------------|-------------------------------|---------|
| Cerebellum              | 75 $\pm$ 13 (n=8)                  | 73 $\pm$ 9 (n=11)             | p=0.91  |
| Frontal Pole            | 58 $\pm$ 8 (n=6)                   | 124 $\pm$ 16 (n=10)           | p<0.006 |
| Superior/Middle Frontal | 54 $\pm$ 10 (n=8)                  | 130 $\pm$ 21 (n=11)           | p<0.005 |
| Superior Temporal       | 66 $\pm$ 17 (n=8)                  | 121 $\pm$ 14 (n=11)           | p<0.02  |
| Head of Caudate         | 73 $\pm$ 11 (n=4)                  | 123 $\pm$ 22 (n=7)            | p=0.14  |
| Inferior Parietal       | 76 $\pm$ 13 (n=8)                  | 137 $\pm$ 19 (n=11)           | p<0.03  |
| Occipital               | 64 $\pm$ 16 (n=8)                  | 95 $\pm$ 12 (n=11)            | p=0.15  |

\*The CBR for each sample is obtained by dividing the absolute value of [ $^{14}$ C]Chrysamine G binding in that sample by the absolute value of [ $^{14}$ C]Chrysamine G binding in the cerebellar sample from that same brain. The values in the table are the average CBRs from each brain area ( $\pm$  SEM). High- and low-plaque AD brains combined.

Figure 9A and 9B shows the binding of [ $^{14}$ C]Chrysamine G to six brain areas normalized to the cerebellum of the same brain. The binding of Chrysamine G to AD brain areas in AD brains having more than 20 NPs/x200 magnification, "High Plaque AD Brains", is shown in Figure 9A. The binding of Chrysamine G to AD brain areas in AD brains having less than 20 NPs/x200 magnification, "Low Plaque AD Brains", is shown in Figure 9B. In all brain areas, the binding to AD brain is significantly greater than the binding to control (see Table 3). In superior/middle frontal cortex, there is no overlap between control and any of the AD samples. In all brain areas except the occipital cortex, there is no overlap between control

-72-

and the AD samples having >20 NPs/x200 magnification. In brain areas with the least deposition of classical NPs, such as the occipital cortex (and cerebellum), the greatest overlap between AD and control was observed.

5        Figure 9C shows the data from two patients who had Down's syndrome. Down's syndrome patients all develop deposits of A $\beta$  by their fourth decade and many develop AD. Wisniewski et al., *Neurology* 35: 957 (1985); Schapiro et al., *Neurobiol. Aging* 13, 723 (1992). Both  
10       of these patients showed [<sup>14</sup>C]Chrysamine G binding above the control range. Since the younger patient (23 years old) had amyloid deposits but was not yet clinically demented, Figure 9C suggests that Chrysamine G can detect differences from control in non-demented  
15       patients destined to develop AD long before the dementia is clinically evident.

      The compounds and method of the invention provide two useful measurements for differentiating AD brain from normal brain; either (1) total Chrysamine G  
20       binding (Table 2) or (2) the ratio of Chrysamine G binding in a given brain area to binding in the cerebellum of the same brain (Table 3). These measurements furnish two great advantages for *in vivo* quantitation of AD neuritic plaques. First, by  
25       providing a means to measure total A $\beta$  binding, rather than specific A $\beta$  binding, the instant invention can quantify A $\beta$  deposition without having to expose the subject to a second injection of radioactive material in order to measure non-specific binding. Because of  
30       this, the data are expressed as total binding only. In all of the experiments presented, specific binding data yields even greater differences between AD and control brain.

      Second, variations in brain uptake of Chrysamine G  
35       derivatives will affect the absolute concentration of Chrysamine G in brain. Some mechanism will be necessary, therefore, to account for these variations

-73-

between subjects. Each patient can serve as his/her own control by finding a brain area that shows little A $\beta$  deposition (i.e., an experimental "blank"). Since classical NPs are exceedingly rare in the cerebellum (Joachim et al., *Am. J. Pathol.* 135: 309 (1989)), Chrysamine G binding to the cerebellum was used as a control for each brain studied. The results were expressed in terms of the ratio of Chrysamine G binding in a given brain area to binding in the cerebellum of the same brain (Figure 9 and Table 3).

For the purposes of *in vivo* quantitation of amyloid in AD, the effect of brain atrophy should be considered. Therefore, when using the Chrysamine G and Chrysamine G derivative probes *in vivo* to quantitate amyloid, brain atrophy can be corrected based on MRI volume measurements. MRI volume measurements performed in conjunction with the method of the invention are analogous to those routinely employed in the art. See, Pearlson, G. and Marsh, L. MAGNETIC RESONANCE IMAGING IN PSYCHIATRY in *Annual Review of Psychiatry* (Vol. 12) Oldham, J. et al., eds. p. 347-381 (1993). Therefore a method for determining the total radioactivity per volume of brain area would use the following equation:

$$\frac{\text{total SPECT or PET signal from brain area "A"}}{\text{MRI determined brain volume (excluding CSF) in brain area "A"}}$$

Designating this measurement as the signal/volume for brain area "A" or  $S/V_A$  means that the cerebellar ratio would be expressed as:

$$\text{Ratio}_A = \frac{S/V_A}{S/V_{CB}}$$

where  $S/V_{CB}$  is the signal/volume in the cerebellum of the same subject during the same imaging study. This ratio from any brain area other than cerebellum from a patient suspected of having AD or other pathological

-74-

condition characterized by the deposition of amyloid could then be compared to the normal range of the analogous ratio from the same brain area of a group of age-matched normal control subjects. The ratio of the binding to brain areas with high deposits of neuritic plaques to the cerebellum can be used as the parameter to distinguish Alzheimer from control subjects.

**EXAMPLE 4      THE OCTANOL-WATER PARTITION COEFFICIENTS**  
**OF CHRYSAMINE G, CHRYSAMINE G**  
**DERIVATIVES, AND CONGO RED**

The octanol-water partition coefficient is a measure of the relative lipophilicity of a compound. The more lipophilic a compound, the more likely it is to cross the blood-brain barrier. See, Goodman and Gilman's THE PHARMACOLOGICAL BASIS FOR THERAPEUTICS (7th Ed.). The octanol/water partition coefficient of Chrysamine G is  $60.22 \pm 3.97$  and that of Congo red is  $0.665 \pm 0.037$  ( $p < 0.001$ ). This suggests that Chrysamine G is approximately 90 times more lipophilic than Congo red and therefore is theoretically more likely to cross the mammalian blood-brain barrier. The octanol/water partition coefficients for the 3-iodo and 3,3'-diiodo derivatives of Chrysamine G (Figure 1) are  $72.53 \pm .74$  and  $112.9 \pm 7.3$ , respectively. These octanol/water partition coefficients show that these derivatives, which are non-radioactive analogues of some of the radiolabeled Chrysamine G derivatives to be used for *in vivo* studies, are up to 170 times more lipophilic than Congo red and up to twice as lipophilic as Chrysamine G. This suggests they will enter the brain much better than either Congo red or Chrysamine G.

-75-

**EXAMPLE 5      THE ABILITY OF CHRYSAMINE G AND  
CHRYSAMINE G DERIVATIVES TO CROSS THE  
BLOOD-BRAIN BARRIER AND METABOLISM OF  
CHRYSAMINE G**

5            Use of the amyloid probes to diagnose AD *in vivo* requires them to be able to cross the blood-brain barrier and gain access to parenchymal amyloid deposits.

10           The ability of Chrysamine G to cross the blood-brain barrier was studied in Swiss-Webster mice. After i.v. injection, the brain/blood ratio measured at 15 min was over 10:1 and approached 20:1 by 35 min (Figure 10). The radioactivity in brain stayed nearly constant over this period, but decreased in the blood and  
15           increased in the liver. The brain/kidney ratio was highest at 15 min (over the time points sampled) and approached 0.5. When brain and liver were extracted 60 min after i.v. injection of [<sup>14</sup>C]Chrysamine G, >95% of the recovered radioactivity co-eluted with authentic  
20           Chrysamine G on reverse phase HPLC, indicating no significant metabolism of Chrysamine G over this period of time.

            Chrysamine G does get into normal mouse brain, and the brain/blood ratio is high. The radioactivity  
25           in brain remained relatively constant over the first 30 min while decreasing in blood and increasing in liver. This suggests that the high brain/blood ratio is more a result of efficient removal of Chrysamine G from the blood by the liver than to further accumulation in the  
30           brain. At 60 min, essentially all of the radioactivity found in the brain and liver proved to be unchanged Chrysamine G. Congo red does not cross the blood-brain barrier well. Tubis et al., *J. Amer. Pharm. Assn.* 49: 422 (1960). Most of the Congo red is cleared by the  
35           liver and spleen and the brain/kidney ratio achieved in guinea pigs is approximately 0.07. Tubis et al., *supra*. Chrysamine G also is cleared by the liver, but

-76-

has greater entry into the brain.

In vivo animal testing provides yet a further basis for determining dosage ranges, efficacy of transfer through the blood barrier and binding ability.

5 Particularly preferred for this purpose are the transgenic mouse model of Games et al., (*Nature* 373: 523 (1995)) or Hsiao et al. *Science* 274: 99-102 (1996) and the "senile animal" model for cerebral amyloidosis; i.e., animals such as the transgenic mice or aged dogs  
10 or monkeys, which are known to develop variable numbers of Alzheimer-type cerebral neuritic plaques, see Wisniewski et al., *J. Neuropathol. & Exp. Neurol.* 32: 566 (1973), Selkoe et al., *Science* 235: 873 (1987), are tested for binding and detection efficacy. This in  
15 vivo assay requires control-biopsy or necropsy monitoring to confirm and quantify the presence of amyloid deposits.

Other suitable animal models for use in testing the compositions and methods of the present invention  
20 are produced transgenically. For instance, Quon et al., *Nature*, 352: 239-241 (1991) used rat neural-specific enolase promoter inhibitor domain to prepare transgenic mice. See also, Wirak et al., *Science*, 253: 323-325 (1991). Still other models have been produced  
25 by Intracranial administration of the  $\beta$ /A4 peptide directly to animals (Tate et al., *Bull. Clin. Neurosci.*, 56: 131-139 (1991)).

It is noted that none of the in vivo animal models may turn out to be extremely good models for AD  
30 neuropathology. Instead, they may more closely model the amyloid deposition of normal aging. This is particularly true of the aged-mammal models. All of these models show a preponderance of diffuse plaques as discussed above for the aged dog model. While there is  
35 some cerebrovascular amyloid, there are few neuritic plaques, except in the Games et al. and Hsiao et al. transgenic mouse models. The other transgenic mouse

-77-

models often show only diffuse plaques. Therefore, while these models may be useful for studying distribution of the probes in the brain, there is a fairly low probability that these models would show the same quantitative differences that would be expected to be seen in AD brain based on the *in vitro* studies of Chrysamine G binding to AD brain described above.

**Evaluating the Ability of Alkyl, Alkenyl or Alkynyl Chrysamine G Derivatives to Cross the Human Blood-Brain Barrier**

A dose of approximately 10 mCi of an appropriately radiolabeled derivative of Chrysamine G with a specific activity of approximately 500 Ci/mole or higher is injected intravenously into normal subjects and patients suspected of having AD and monitored by SPECT or PET imaging to analyze the detectability of the derivative in brain relative to other organs and to define the time course of detectability in the brain. A dose which can be reliably detected is defined as a "imaging effective dose."

**Evaluation of Alkyl, Alkenyl or Alkynyl Chrysamine G Derivatives to Distinguish AD from Age-Matched Controls in Humans**

An imaging-effective dose of an appropriately, radioactively labeled derivative of Chrysamine G is injected into a subject suspected of having brain amyloid deposition due to pathological conditions such as AD. After a period of 15 minutes to 24 hours, the radioactive signal from brain is detected by SPECT or PET. Radioactivity is simultaneously detected in all brain areas included in the field of view of the detector. This field of view will be set up so as to include large portions of the cerebellum, superior temporal cortex, superior/middle frontal cortex, and intervening brain regions. An MRI scan will be performed prior to the study so that corrections can be

-78-

made for brain atrophy in the areas of interest by methods discussed in Example 3. The  $S/V_A$ ,  $S/V_{CB}$ , and  $Ratio_A$  variables discussed in Example 3 will be calculated and compared to analogous normative ratios obtained previously from age-matched normal control subjects.

**EXAMPLE 6      HISTOLOGIC LOCALIZATION OF THE BINDING OF AN ALKENYL CHRYSAMINE G DERIVATIVE TO AMYLOID**

10        The top frame of Figure 11 demonstrates a human AD brain stained by 1,4-bis(2-(3-carboxy-4-hydroxyphenyl)ethen-1-yl)-benzene. The staining method was that of Stokes and Trickey, *J. Clin. Pathol.* 26: 241-242 (1973) with 1,4-bis(2-(3-carboxy-4-hydroxyphenyl)ethen-1-yl)-benzene substituted for Congo red. This staining is much more intense than that observed with Chrysamine G, Congo red, or Thioflavin S. Numerous amyloid plaques and neuropil threads can be readily identified as well as a neurofibrillary tangle.

15        The bottom photomicrograph in Figure 11 shows a section of transgenic mouse brain [Tg(HuAPP695.SWE)2576; Hsiao et al. *Science* 274: 99-102 (1996)] stained with 1,4-bis(2-(3-carboxy-4-hydroxyphenyl)ethen-1-yl)-benzene. The dense amyloid plaques is intensely stained.

20        Cerebrovascular amyloid also is intensely stained in both human and mouse brain (data not shown).



-79-

**EXAMPLE 7      ASSESSING THE TOXICITY OF ALKYL, ALKENYL**  
**OR ALKYNYL CHRYSAMINE G DERIVATIVES**

At doses of 10 and 100 mg/kg non-radioactive Chrysamine G administered intraperitoneally, no notable behavioral effects or toxicity were observed in mice for periods up to 72 hrs. The doses of [<sup>14</sup>C]Chrysamine G administered were on the order of 1 mg/kg.

Chrysamine G appeared to show little acute toxicity based on attempts to establish an LD<sub>50</sub>. Even when the maximum volume that can be injected into a mouse without harming it just from fluid volume effects (approx. 0.025 ml/g) of a saturated solution of Chrysamine G was injected into mice (100 mg/kg), there were no behavioral changes noted for at least 72 hrs, the longest period tested. Doses required for detection of radiolabeled derivatives by SPECT or PET would be orders of magnitude below this dose.

Congo red has been safely injected into humans in quantities much greater than would be used for the radioactive Chrysamine G derivatives. The LD<sub>50</sub> for Congo red has been shown to be 190 mg/kg mouse (Tubis et al., *J. Amer. Pharm. Assoc.* 49: 422 (1960)), which is similar to the >100 mg/kg LD<sub>50</sub> shown for Chrysamine G. Thus, these two chemically similar compounds cause similar low toxicities in mice.

Other alkyl, alkenyl or alkynyl Chrysamine G derivatives can similarly be tested for toxicity in mice and other higher mammals by injecting a wide range of concentrations and studying the animals for various signs of toxicity by methods well known in the art. See, Goodman and Gilman's *THE PHARMACOLOGICAL BASIS FOR THERAPEUTICS* (7th Ed.).

-80-

**EXAMPLE 8      ASSESSING THE ABILITY OF CHRYSAMINE G TO**  
**PROTECT AGAINST A $\beta$  (25-35)-INDUCED**  
**TOXICITY**

5      **Protection from A $\beta$  (25-35)-induced Toxicity in PC-12 cells**

Rat pheochromocytoma cells (PC-12) were grown in RPMI 1640 media with 10% fetal bovine serum. Approximately 5,000 exponentially growing cells were plated in 96-well plates in a volume of 100  $\mu$ l of media and allowed to incubate at 37°C overnight. The A $\beta$  (25-35), which had been pre-aggregated at 37°C for 7 days, was pre-incubated with Chrysamine G (CG) or related compounds in aqueous solution prior to addition of 20  $\mu$ l to achieve the final concentrations given (0.01 to 10  $\mu$ M A $\beta$ (25-35) and 0.03 to 20  $\mu$ M CG). The cells were incubated for 24 hrs prior to the addition of 13.3  $\mu$ l of 5 mg/ml MTT (3, (4,5-dimethylthiazol-2-yl)2,5-diphenyltetrazolium bromide) in sterile phosphate buffered saline. After 4.5 hrs at 37°C, 100  $\mu$ l of extraction buffer (20% w/v SDS in 50% DMF/water; pH adjusted to 4.7 with 2.5% of 80% acetic acid and 2.5% 1N HCl) was added and the plates were incubated overnight. Hansen et al., *J. Immunol. Methods* 119: 203 (1989). Color development was then measured at 560 nm. Maximum viability was defined as the absorbance of control wells to which only the 20  $\mu$ l of distilled, deionized H<sub>2</sub>O was added. Maximum toxicity was defined by wells in which the cells were lysed by addition of 0.1% (final concentration) of Triton X-100.

30      Incubation of PC12 cells with A $\beta$ (25-35) results in a concentration-dependent decrease in the ability of these cells to reduce MTT (Figure 12). Figure 12 shows the effect of increasing concentrations of A $\beta$ (25-35) in the presence and absence of Chrysamine G on the cellular redox activity of PC12 cells as measured by MTT reduction. The reduction product of MTT absorbs at 560 nm which is plotted on the vertical axis. The

-81-

effect of A $\beta$ (25-35) alone is shown in the filled bars and shows a dose dependent decrease in MTT reduction. Significant differences from control (no A $\beta$ , no Chrysamine G) are shown in white numbers inside the filled bars. The protective effect of 20  $\mu$ M Chrysamine G is shown in the open bars. Significant differences between MTT reduction in the presence and absence of Chrysamine G are shown in black numbers inside the open bars.

Figure 13 demonstrates the protective effect of increasing concentrations of Chrysamine G against the A $\beta$ (25-35)-induced reduction of cellular redox activity of PC12 cells. The effect of Chrysamine G in the absence of A $\beta$ (25-35) is shown in the filled bars. There was no significant difference between control (no A $\beta$ , no Chrysamine G) and any of the concentrations of Chrysamine G in the absence of A $\beta$ (25-35). MTT reduction in the presence of 1  $\mu$ M A $\beta$ (25-35) and increasing concentrations of Chrysamine G is shown in the open bars. Significant differences in MTT reduction between the presence and absence of A $\beta$ (25-35) at each concentration of Chrysamine G are shown in white numbers inside the filled bars. Significant differences in MTT reduction between the A $\beta$ (25-35) control (no Chrysamine G) and A $\beta$ (25-35) plus increasing concentrations of Chrysamine G are shown in black numbers inside the open bars.

As has previously been reported, Congo red protects against the A $\beta$ -induced toxicity at concentrations over 2  $\mu$ M, achieving complete protection by 20  $\mu$ M. Burgevin et al. *NeuroReport* 5: 2429 (1994); Lorenzo and Yankner, *Proc. Natl. Acad. Sci.* 91: 12243 (1994); Pollack et al., *Neuroscience Letters* 184: 113 (1995); Pollack et al. *Neuroscience Letters* 197: 211 (1995). Chrysamine G shows a protective effect which is dependent on both the concentration of A $\beta$ (25-35) (Figure 12) as well as the concentration of Chrysamine

-82-

G (Figure 13). The protective effect of Chrysamine G is evident at 0.2  $\mu$ M, a concentration very close to the  $K_i$  of Chrysamine G for binding to synthetic A $\beta$ , 0.37  $\mu$ M (Figure 1). Chrysamine G appears to be more potent than Congo red, showing effects in the range of 0.1 to 1.0  $\mu$ M. This is consistent with the  $K_i$  values for binding to synthetic A $\beta$  of 0.37  $\mu$ M for Chrysamine G and 2.8  $\mu$ M for Congo red (Figure 1).

In another experiment (Figure 14), the effect of Chrysamine G and the phenol derivative (see Figure 1), which does not bind A $\beta$ , was examined in cells incubated with 1  $\mu$ M A $\beta$ (25-35). Chrysamine G showed protective effects at 0.1 and 1  $\mu$ M, but the phenol derivative showed no protective effects, and perhaps enhanced the toxicity of A $\beta$ .

These results suggest that the lipophilic derivative of Congo red, Chrysamine G, prevents A $\beta$ -induced cytotoxicity in cell culture at concentrations very similar to those at which it binds A $\beta$ . This protection shows structural specificity since the phenol derivative which does not bind to synthetic A $\beta$  also does not prevent A $\beta$ -induced cytotoxicity. Since Chrysamine G partitions into the brain well, these results provide evidence that Chrysamine G and A $\beta$ -binding derivatives of Chrysamine G have therapeutic potential in the treatment of AD.

The mechanism of the protective effect of Chrysamine G is unknown at present. Two broad possibilities exist. First, Chrysamine G could interfere with the aggregation of A $\beta$ . Second, Chrysamine G could interfere with the effects (direct or indirect) of A $\beta$  on the target cells. Congo red does inhibit aggregation of A $\beta$  as well as protect against the toxic effects of aggregated A $\beta$ . Lorenzo and Yankner, *Proc. Natl. Acad. Sci.* 91: 12243 (1994). Interference with aggregation is unlikely in the above experiment since the A $\beta$  was pre-aggregated prior to


-83-

incubation with Chrysamine G. Thus, inhibition of aggregation could prove to be an important therapeutic effect of Chrysamine G but is not a likely explanation for the protective effects of Chrysamine G against pre-  
5 aggregated A $\beta$ . The model of Chrysamine G binding to A $\beta$  described in Figure 7, displays how Chrysamine G could "coat" the surface of A $\beta$ . This may change how the fibrillar deposits are recognized by cell-surface  
10 receptors or other macromolecules such as complement proteins and interfere with the toxic effects of A $\beta$  which may be mediated by these macromolecules. It is likely that Chrysamine G and Congo red exert multiple effects, both before and after the aggregation of A $\beta$ .  
15 This is advantageous from a therapeutic point of view since patients are likely to present at a time when there are pre-existing A $\beta$  aggregates as well as ongoing amyloid deposition.

-84-

What Is Claimed Is:

1. An amyloid binding compound of Formula I or a water soluble, non-toxic salt thereof:



wherein:


$Z$  is either,  $C \equiv C$ ,  $CR' = CR'$ ,  $CR'_2 - CR'_2$ ,  $C \equiv C - C \equiv C$ ,  $CR' = CR' - CR' = CR'$ ,  $C \equiv C - CR' = CR'$ , or  $CR'_2 - CR'_2 - CR'_2$  (where each  $R'$  independently represents H or a lower alkyl group);

$X$  is  $C(R'')_2$

(wherein each  $R''$  independently is H, F, Cl, Br, I, a lower alkyl group,  $(CH_2)_nOR'$  where  $n=1, 2$ , or  $3$ ,  $CF_3$ ,  $CH_2-CH_2F$ ,  $O-CH_2-CH_2F$ ,  $CH_2-CH_2-CH_2F$ ,  $O-CH_2-CH_2-CH_2F$ , CN,  $(C=O)-R'$ ,  $N(R')_2$ ,  $NO_2$ ,  $(C=O)N(R')_2$ ,  $O(CO)R'$ ,  $OR'$ ,  $SR'$ ,  $COOR'$ ,  $R_{Ph}$ ,  $CR' = CR' - R_{Ph}$ ,  $CR'_2 - CR'_2 - R_{Ph}$  (where  $R_{Ph}$  represents an unsubstituted or substituted phenyl group with the phenyl substituents being chosen from any of the non-phenyl substituents defined for  $R''$ , a tri-alkyl tin, a tetrazole or oxadiazole of the

-85-


form:




wherein R' is H or a lower alkyl group)

or X is CR'=CR', N=N, C=O, O, NR' (where R' represents H or a lower alkyl group), S, or SO<sub>2</sub>;


each R<sub>1</sub> and R<sub>2</sub> independently is H, F, Cl, Br, I, a lower alkyl group, (CH<sub>2</sub>)<sub>n</sub>OR' where n=1, 2, or 3, CF<sub>3</sub>, CH<sub>2</sub>-CH<sub>2</sub>F, O-CH<sub>2</sub>-CH<sub>2</sub>F, CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>F, O-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>F, CN, (C=O)-R', N(R')<sub>2</sub>, NO<sub>2</sub>, (C=O)N(R')<sub>2</sub>, O(CO)R', OR', SR', COOR', a tri-alkyl tin, R<sub>ph</sub>, CR'=CR'-R<sub>ph</sub>, CR'<sub>2</sub>-CR'<sub>2</sub>-R<sub>ph</sub> (where R<sub>ph</sub> represents an unsubstituted or substituted phenyl group with the phenyl substituents being chosen from any of the non-phenyl substituents defined for R<sub>1</sub> and R<sub>2</sub>, a tetrazole or oxadiazole of the form:



(wherein R' is H or a lower alkyl group), or a triazene of the form:

(wherein R<sub>8</sub> and R<sub>9</sub> are lower alkyl groups)

or




-86-

each Q is independently selected from one of the following structures:

IA, IB, IC, ID, IE, IF and IG, wherein


IA has the following structure:



wherein:

each of R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, or R<sub>7</sub> independently is defined the same as R<sub>1</sub> above;


IB has the following structure:



wherein:

each of R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub>, R<sub>14</sub>, R<sub>15</sub>, or R<sub>16</sub> independently is defined the same as R<sub>1</sub> above;

IC has the following structure:






-87-

wherein:

each of  $R_{17}$ ,  $R_{18}$ ,  $R_{19}$ ,  $R_{20}$ , or  $R_{21}$  independently is defined the same as  $R_1$  above;


ID has the following structure:




wherein:

each of  $R_{22}$ ,  $R_{23}$ , or  $R_{24}$  independently is defined the same as  $R_1$  above

and




represents a heterocyclic ring of one of the six following formulas:



IE has the following structure:


-88-




wherein:

each of  $R_{25}$ ,  $R_{26}$ , or  $R_{27}$  independently is defined the same as  $R_1$  above

and




represents a heterocyclic ring of one of the six following formulas:



IF has the following structure:

-89-




wherein:

exactly one of  $R_{28}$ ,  $R_{29}$ ,  $R_{30}$ ,  $R_{31}$  or  $R_{32}$  is the  $\text{Z}$  link defined for Formula I above and each other  $R_{28}$ ,

$R_{29}$ ,  $R_{30}$ ,  $R_{31}$  or  $R_{32}$  independently is defined the same as  $R_1$  above;

IG has the following structure:

(IG)



wherein:

exactly one of  $R_{33}$ ,  $R_{34}$ ,  $R_{35}$ ,  $R_{36}$ ,  $R_{37}$ ,  $R_{38}$ , or  $R_{39}$  is the

$\text{Z}$  link defined for Formula I above and each other

-90-

$R_{33}$ ,  $R_{34}$ ,  $R_{35}$ ,  $R_{36}$ ,  $R_{37}$ ,  $R_{38}$ , or  $R_{39}$  independently is defined the same as  $R_1$  above.

2. The compound of claim 1, wherein at least one of the substituents  $R_1$ - $R_7$  and  $R_{10}$ - $R_{39}$  is selected from the group consisting of  $^{131}\text{I}$ ,  $^{123}\text{I}$ ,  $^{76}\text{Br}$ ,  $^{75}\text{Br}$ ,  $^{18}\text{F}$ ,  $\text{CH}_2\text{-CH}_2\text{-}^{18}\text{F}$ ,  $\text{O-CH}_2\text{-CH}_2\text{-}^{18}\text{F}$ ,  $\text{CH}_2\text{-CH}_2\text{-CH}_2\text{-}^{18}\text{F}$ ,  $\text{O-CH}_2\text{-CH}_2\text{-CH}_2\text{-}^{18}\text{F}$ ,  $^{19}\text{F}$ ,  $^{125}\text{I}$ , and a carbon-containing substituent as specified in Formula I wherein at least one carbon is  $^{11}\text{C}$  or  $^{13}\text{C}$ .

3. The compound of claim 1, wherein said compound binds to A $\beta$  with a dissociation constant ( $K_D$ ) between 0.0001 and 10.0  $\mu\text{M}$  when measured by binding to synthetic A $\beta$  peptide or Alzheimer's Disease brain tissue.

4. A method for synthesizing a compound of claim 1, wherein at least one of the substituents  $R_1$ - $R_7$  and  $R_{10}$ - $R_{39}$  is selected from the group consisting of  $^{131}\text{I}$ ,  $^{125}\text{I}$ ,  $^{123}\text{I}$ ,  $^{76}\text{Br}$ ,  $^{75}\text{Br}$ ,  $^{18}\text{F}$  and  $^{19}\text{F}$ , comprising the step of reacting a compound of claim 1, wherein at least one of the substituents  $R_1$ - $R_7$  and  $R_{10}$ - $R_{39}$  is a tri-alkyl tin or a triazene, with a halogenating agent containing  $^{131}\text{I}$ ,  $^{125}\text{I}$ ,  $^{123}\text{I}$ ,  $^{76}\text{Br}$ ,  $^{75}\text{Br}$ ,  $^{18}\text{F}$  or  $^{19}\text{F}$ .

5. A pharmaceutical composition for *in vivo* imaging of amyloid deposits, comprising (a) a compound of claim 2, and (b) a pharmaceutically acceptable carrier.

6. An *in vivo* method for detecting amyloid deposits in a subject, comprising the steps of:

(a) administering a detectable quantity of the pharmaceutical composition of claim 5, and

(b) detecting the binding of said compound to amyloid deposit in said subject.

7. The method of claim 6, wherein said amyloid

-91-

deposit is located in the brain of a subject.

8. The method of claim 6, wherein said subject is suspected of having a disease or syndrome selected from the group consisting of Alzheimer's Disease, familial Alzheimer's Disease, Down's Syndrome and homozygotes for the apolipoprotein E4 allele.

9. The method of claim 6, wherein said detecting is selected from the group consisting of gamma imaging, magnetic resonance imaging and magnetic resonance spectroscopy.

10. The method of claim 9, wherein said gamma imaging is either PET or SPECT.

11. The method of claim 7, wherein said pharmaceutical composition is administered by intravenous injection.


12. The method of claim 7, wherein the ratio of (i) binding of said compound to a brain area other than the cerebellum to (ii) binding of said compound to the cerebellum, in said subject, is compared to said ratio in normal subjects.

13. A method of inhibiting cell degeneration and toxicity associated with fibril formation in an amyloidosis-associated condition, said method comprising the step of administering to a subject having or suspected of having such condition, a pharmaceutically effective amount of Chrysamine G or a derivative thereof.

14. The method of claim 13, wherein said derivative of Chrysamine G is an amyloid binding compound of Formula

-92-

I or a water soluble, non-toxic salt thereof:



wherein:


Z is either,  $C \equiv C$ ,  $CR' = CR'$ ,  $CR'_2 - CR'_2$ ,

$C \equiv C - C \equiv C$ ,  $CR' = CR' - CR' = CR'$ ,  $C \equiv C - CR' = CR'$ , or  $CR'_2 - CR'_2 - CR'_2$  (where each R' independently represents H or a lower alkyl group);

X is  $C(R'')_2$

(wherein R'' is each independently is H, F, Cl, Br, I, a lower alkyl group,  $(CH_2)_nOR'$  where n=1, 2, or 3,  $CF_3$ ,  $CH_2-CH_2F$ ,  $O-CH_2-CH_2F$ ,  $CH_2-CH_2-CH_2F$ ,  $O-CH_2-CH_2-CH_2F$ ,  $CN$ ,  $(C=O)-R'$ ,  $N(R')_2$ ,  $NO_2$ ,  $(C=O)N(R')_2$ ,  $O(CO)R'$ ,  $OR'$ ,  $SR'$ ,  $COOR'$ ,  $R_{ph}$ ,  $CR' = CR' - R_{ph}$ ,  $CR'_2 - CR'_2 - R_{ph}$  (where  $R_{ph}$  represents an unsubstituted or substituted phenyl group with the phenyl substituents being chosen from any of the non-phenyl substituents defined for R'', a tri-alkyl tin, a tetrazole or oxadiazole of the form:


-93-




wherein R' is H or a lower alkyl group)

or X is CR'=CR', N=N, C=O, O, NR' (where R' represents H or a lower alkyl group), S, or SO<sub>2</sub>;

each R<sub>1</sub> and R<sub>2</sub> independently is H, F, Cl, Br, I, a lower alkyl group, (CH<sub>2</sub>)<sub>n</sub>OR' where n=1, 2, or 3, CF<sub>3</sub>, CH<sub>2</sub>-CH<sub>2</sub>F, O-CH<sub>2</sub>-CH<sub>2</sub>F, CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>F, O-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>F, CN, (C=O)-R', N(R')<sub>2</sub>, NO<sub>2</sub>, (C=O)N(R')<sub>2</sub>, O(CO)R', OR', SR', COOR', a tri-alkyl tin, R<sub>Ph</sub>, CR'=CR'-R<sub>Ph</sub>, CR'<sub>2</sub>-CR'<sub>2</sub>-R<sub>Ph</sub> (where R<sub>Ph</sub> represents an unsubstituted or substituted phenyl group with the phenyl substituents being chosen from any of the non-phenyl substituents defined for R<sub>1</sub> and R<sub>2</sub>, a tetrazole or oxadiazole of the form:




(wherein R' is H or a lower alkyl group), or a triazene of the form:



(wherein R<sub>8</sub> and R<sub>9</sub> are lower alkyl groups)

or




-94-

each Q is independently selected from one of the following structures:

IA, IB, IC, ID, IE, IF and IG, wherein


IA has the following structure:



wherein:

each of R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, or R<sub>7</sub> independently is defined the same as R<sub>1</sub> above;


IB has the following structure:



wherein:

each of R<sub>10</sub>, R<sub>11</sub>, R<sub>12</sub>, R<sub>13</sub>, R<sub>14</sub>, R<sub>15</sub>, or R<sub>16</sub> independently is defined the same as R<sub>1</sub> above;

IC has the following structure:



wherein:


each of R<sub>17</sub>, R<sub>18</sub>, R<sub>19</sub>, R<sub>20</sub>, or R<sub>21</sub> independently is



-95-

defined the same as  $R_1$  above;


ID has the following structure:




wherein:

each of  $R_{22}$ ,  $R_{23}$ , or  $R_{24}$  independently is defined the same as  $R_1$  above

and




represents a heterocyclic ring of one of the six following formulas:



IE has the following structure:


-96-




wherein:

each of  $R_{25}$ ,  $R_{26}$ , or  $R_{27}$  independently is defined the same as  $R_1$  above

and




represents a heterocyclic ring of one of the six following formulas:



IF has the following structure:


-97-



wherein:

exactly one of  $R_{28}$ ,  $R_{29}$ ,  $R_{30}$ ,  $R_{31}$  or  $R_{32}$  is the  $\text{Z}$  link defined for Formula I above and each other  $R_{28}$ ,  $R_{29}$ ,  $R_{30}$ ,  $R_{31}$  or  $R_{32}$  independently is defined the same as  $R_1$  above;

IG has the following structure:



wherein:

exactly one of  $R_{33}$ ,  $R_{34}$ ,  $R_{35}$ ,  $R_{36}$ ,  $R_{37}$ ,  $R_{38}$ , or  $R_{39}$  is the

$\text{Z}$  link defined for Formula I above and each other

-98-

$R_{33}$ ,  $R_{34}$ ,  $R_{35}$ ,  $R_{36}$ ,  $R_{37}$ ,  $R_{38}$ , or  $R_{39}$  independently is defined the same as  $R_1$  above.

15. The method of claim 13, wherein said amyloidosis-associated condition is selected from the group consisting of Alzheimer's Disease, Down's Syndrome, Type 2 diabetes mellitus, hereditary cerebral hemorrhage amyloidosis (Dutch), amyloid A (reactive), secondary amyloidosis, familial mediterranean fever, familial amyloid nephropathy with urticaria and deafness (Muckle-wells Syndrome), amyloid lambda L-chain or amyloid kappa L-chain (idiopathic, myeloma or macroglobulinemia-associated) A beta 2M (chronic hemodialysis), ATTR (familial amyloid polyneuropathy (Portuguese, Japanese, Swedish), familial amyloid cardiomyopathy (Danish), isolated cardiac amyloid, (systemic senile amyloidosises), AIAPP or amylin insulinoma, atrial naturetic factor (isolated atrial amyloid), procalcitonin (medullary carcinoma of the thyroid), gelsolin (familial amyloidosis (Finnish)), cystatin C (hereditary cerebral hemorrhage with amyloidosis (Icelandic)), AApo-A-I (familial amyloidotic polyneuropathy - Iowa), AApo-A-II (accelerated senescence in mice), fibrinogen-associated amyloid; and Asor or Pr P-27 (scrapie, Creutzfeld Jacob disease, Gertsman-Straussler-Scheinker syndrome, bovine spongiform encephalitis) or in cases of persons who are homozygous for the apolipoprotein E4 allele, and the condition associated with homozygosity for the apolipoprotein E4 allele.

16. A pharmaceutical composition for the prevention of cell degeneration and toxicity associated with fibril formation in amyloidosis-associated conditions comprising a Chrysamine G derivative of claim 1 and a

-99-

pharmaceutically acceptable carrier.

17. A method of detecting amyloid deposits in biopsy or post-mortem human or animal tissue comprising the steps of:

(a) incubating formalin-fixed tissue with a solution of a compound of claim 1 to form a labelled deposit and then,

(b) detecting the labelled deposits.

18. The method of claim 17 wherein said solution is composed of 25-100% ethanol (the remainder being water) saturated with a compound of claim 1.

19. The method of claim 17 wherein said detecting is effected by microscopic techniques selected from the group consisting of bright-field, fluorescence, laser-confocal, and cross-polarization microscopy.


20. A method of quantifying the amount of amyloid in biopsy or post-mortem tissue comprising the steps of:

- a) incubating a radiolabeled derivative of Chrysamine G with a homogenate of biopsy or post-mortem tissue,
- b) separating the tissue-bound from the tissue-unbound radiolabeled Chrysamine G derivative,
- c) quantifying the tissue-bound radiolabeled Chrysamine G derivative, and
- d) converting the units of tissue-bound radiolabeled Chrysamine G derivative to units of micrograms of amyloid per 100 mg of tissue by comparison with a standard.

21. The method of claim 20, wherein said radiolabeled derivative of Chrysamine G is an amyloid

-100-

binding compound of Formula I or a water soluble, non-toxic salt thereof:



wherein:


$Z$  is either,  $C\equiv C$ ,  $CR'=CR'$ ,  $CR'_2-CR'_2$ ,

$C\equiv C-C\equiv C$ ,  $CR'=CR'-CR'=CR'$ ,  $C\equiv C-CR'=CR'$ , or  $CR'_2-CR'_2-CR'_2$  (where each  $R'$  independently represents H or a lower alkyl group);

$X$  is  $C(R'')_2$

(wherein  $R''$  is each independently is H, F, Cl, Br, I, a lower alkyl group,  $(CH_2)_nOR'$  where  $n=1, 2$ , or  $3$ ,  $CF_3$ ,  $CH_2-CH_2F$ ,  $O-CH_2-CH_2F$ ,  $CH_2-CH_2-CH_2F$ ,  $O-CH_2-CH_2-CH_2F$ ,  $CN$ ,  $(C=O)-R'$ ,  $N(R')_2$ ,  $NO_2$ ,  $(C=O)N(R')_2$ ,  $O(CO)R'$ ,  $OR'$ ,  $SR'$ ,  $COOR'$ ,  $R_{Ph}$ ,  $CR'=CR'-R_{Ph}$ ,  $CR'_2-CR'_2-R_{Ph}$  (where  $R_{Ph}$  represents an unsubstituted or substituted phenyl group with the phenyl substituents being chosen from any of the non-phenyl substituents defined for  $R''$ , a tri-alkyl tin, a tetrazole or oxadiazole of the form:


-101-




wherein R' is H or a lower alkyl group)

or X is CR'=CR', N=N, C=O, O, NR' (where R' represents H or a lower alkyl group), S, or SO<sub>2</sub>;

each R<sub>1</sub> and R<sub>2</sub> independently is H, F, Cl, Br, I, a lower alkyl group, (CH<sub>2</sub>)<sub>n</sub>OR' where n=1, 2, or 3, CF<sub>3</sub>, CH<sub>2</sub>-CH<sub>2</sub>F, O-CH<sub>2</sub>-CH<sub>2</sub>F, CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>F, O-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>F, CN, (C=O)-R', N(R')<sub>2</sub>, NO<sub>2</sub>, (C=O)N(R')<sub>2</sub>, O(CO)R', OR', SR', COOR', a tri-alkyl tin, R<sub>ph</sub>, CR'=CR'-R<sub>ph</sub>, CR'<sub>2</sub>-CR'<sub>2</sub>-R<sub>ph</sub> (where R<sub>ph</sub> represents an unsubstituted or substituted phenyl group with the phenyl substituents being chosen from any of the non-phenyl substituents defined for R<sub>1</sub> and R<sub>2</sub>, a tetrazole or oxadiazole of the form:




(wherein R' is H or a lower alkyl group), or a triazene of the form:



(wherein R<sub>8</sub> and R<sub>9</sub> are lower alkyl groups)

or




-102-

each Q is independently selected from one of the following structures:

IA, IB, IC, ID, IE, IF and IG, wherein


IA has the following structure:



wherein:

each of  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$ , or  $R_7$  independently is defined the same as  $R_1$  above;


IB has the following structure:



wherein:

each of  $R_{10}$ ,  $R_{11}$ ,  $R_{12}$ ,  $R_{13}$ ,  $R_{14}$ ,  $R_{15}$ , or  $R_{16}$  independently is defined the same as  $R_1$  above;

IC has the following structure:



wherein:


each of  $R_{17}$ ,  $R_{18}$ ,  $R_{19}$ ,  $R_{20}$ , or  $R_{21}$  independently is



-103-

defined the same as  $R_1$  above;


ID has the following structure:




wherein:

each of  $R_{22}$ ,  $R_{23}$ , or  $R_{24}$  independently is defined the same as  $R_1$  above

and




represents a heterocyclic ring of one of the six following formulas:



IE has the following structure:


-104-




wherein:

each of  $R_{25}$ ,  $R_{26}$ , or  $R_{27}$  independently is defined the same as  $R_1$  above

and




represents a heterocyclic ring of one of the six following formulas:



IF has the following structure:


-105-



wherein:

exactly one of  $R_{28}$ ,  $R_{29}$ ,  $R_{30}$ ,  $R_{31}$  or  $R_{32}$  is the  $\text{Z}$  link defined for Formula I above and each other  $R_{28}$ ,  $R_{29}$ ,  $R_{30}$ ,  $R_{31}$  or  $R_{32}$  independently is defined the same as  $R_1$  above;

IG has the following structure:



wherein:

exactly one of  $R_{33}$ ,  $R_{34}$ ,  $R_{35}$ ,  $R_{36}$ ,  $R_{37}$ ,  $R_{38}$ , or  $R_{39}$  is the  $\text{Z}$  link defined for Formula I above and each other

-106-

$R_{33}$ ,  $R_{34}$ ,  $R_{35}$ ,  $R_{36}$ ,  $R_{37}$ ,  $R_{38}$ , or  $R_{39}$  independently is defined the same as  $R_1$  above.


22. The method of claim 21 wherein at least one of the substituents  $R_1$ - $R_7$  and  $R_{10}$ - $R_{39}$  is labeled with a radiolabel selected from the group consisting of  $^{125}\text{I}$ ,  $^3\text{H}$ , and a carbon-containing substituent as specified in Formula I, wherein at least one carbon is  $^{14}\text{C}$ .

23. A method of distinguishing an Alzheimer's disease brain from a normal brain comprising the steps of:

- a) obtaining tissue from (i) the cerebellum and (ii) another area of the same brain other than the cerebellum, from normal subjects and from subjects suspected of having Alzheimer's disease;
- b) incubating said tissues with a radiolabeled Chrysamine G derivative so that amyloid in said tissue binds with said radiolabeled Chrysamine G derivative;
- c) quantifying the amount of amyloid bound to said radiolabeled Chrysamine G derivative, according to the method of claim 20;
- d) calculating the ratio of the amount of amyloid in the area of the brain other than the cerebellum to the amount of amyloid in the cerebellum;
- e) comparing said ratio for amount of amyloid in said tissue from normal subjects with ratio for amount of amyloid in tissue from subjects suspected of having Alzheimer's disease; and
- f) determining the presence of Alzheimer's disease if said ratio from the brain of a subject suspected of having Alzheimer's disease is above 90% of the ratios obtained from the brains of normal subjects.


1/15


FIG. 1A




2/15

FIG. 1B




**FIG. 2A**


CHRYSAMINE G OR  
SALICYLIC ACID DERIVATIVE.

**FIG. 2B**


CHRYSAMINE G WITH  
GENERAL TRI-ALKYL TIN.

**FIG. 2C**


3-HYDROXY-1,2-  
BENZISOXAZOLE DERIVATIVE.

**FIG. 2D**


PHTHALIMIDE OR  
ISOINDOLE-1,3(2H)-DIONE.

**FIG. 2E**


PHTHALHYDRAZIDE OR 2,3  
BENZODIAZINE-1,4  
(2H,3H)-DIONE DERIV.

**FIG. 2F**


BENZOXAZINE-1,4(3H)-  
DIONE DERIVATIVE.

**FIG. 2G**


(2H)-1,3-BENZOXAZINE-2,4(3H)-  
DIONE DERIVATIVE.

**FIG. 2H**


(3H)-2-BENZAZINE-1,3(2H)-  
DIONE DERIVATIVE.

**FIG. 2I**

1,8-NAPHTHALIMIDE  
DERIVATIVE

**FIG. 2J**


TETRAZOLE DERIVATIVE.

**FIG. 2K**

OXADIAZOLE DERIVATIVE.



4/15

FIG. 3







5/15  
**FIG. 4**



6/15


**FIG. 5**

7/15

**FIG. 6**


8/15


FIG. 7



9/15


FIG. 8



10/15  
**FIG. 9**

11/15

**FIG. 10**




12/15

FIG. 11






13/15


**FIG. 12**

14/15

FIG. 13



15/15

**FIG. 14**

# INTERNATIONAL SEARCH REPORT

International Application No.  
PCT/US 98/07889

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C09B23/14 C07C65/19 C07C65/28 C07C69/88 C07F7/22  
C07B59/00 G01N33/68 A61K31/60 A61K49/00 A61K51/00  
//C07D209/48, C07D261/20, C07D237/32, C07D265/02, C07D265/26,

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C09B C07C C07F C07D C07M C07B G01N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages  | Relevant to claim No. |
|------------|---|-----------------------|
| X          | WO 96 34853 A (UNIVERSITY OF PITTSBURGH) 7 November 1996<br>see whole document, in particular page 74, lines 6-24, and figure 1B, compound 3-(COOH)-C3C | 1-23                  |
| X          | W.E. KLUNK ET AL: NEUROBIOLOGY OF AGING, vol. 15, no. 6, 1994, pages 691-698, XP000602786<br>see figures 2-6; discussion                                | 13                    |
| X          | CH 554 821 A (CIBA-GEIGY AG) 15 October 1974<br>see compounds 134-136, 136b-139, 141-141a, 165-173, 176-179, 184-186, 192-193<br>---<br>-/--            | 1                     |

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

9 September 1998

Date of mailing of the international search report

25/09/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Van Amsterdam, L

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/07889

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D217/24, C07D221/14, C07D257/04, C07D271/10

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages  | Relevant to claim No. |
|------------|---|-----------------------|
| X          | US 3 755 446 A (H. SCHEUERMANN ET AL) 28 August 1973<br>see examples 1-6, 10, 12-13, 18, 23-25, 30-34, 39, 46, 47, 52, 56-57, 61-62, 65, 67-68, 70, 72-73, 75-76<br>--- | 1                     |
| X          | DE 27 36 889 A (CIBA-GEIGY AG) 23 February 1978<br>see examples 1, 4.2-4.4, 5.1, 5.4-5.6, 11.2-11.4<br>---  | 1                     |
| X          | H.H. HOERHOLD ET AL: FASERFORSCH. TEXTILTECH.,<br>vol. 29, no. 6, 1978,<br>pages 393-395, XP002076774<br>see page 393, scheme<br>---                                    | 1                     |

-/--



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

9 September 1998

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Van Amsterdam, L

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 98/07889

| C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT |   |                       |
|--|---|-----------------------|
| Category *   | Citation of document, with indication, where appropriate, of the relevant passages  | Relevant to claim No. |
| X  | <p>DATABASE CHEMABS<br/>           CHEMICAL ABSTRACTS SERVICE, COLUMBUS,<br/>           OHIO, US<br/>           STN, accession no. 113:153324,<br/>           XP002076776<br/>           see abstract; RN 129636-78-8<br/>           -&amp; CHEMICAL ABSTRACTS, vol. 113, no. 18,<br/>           29 October 1990<br/>           Columbus, Ohio, US;<br/>           abstract no. 153324,<br/>           XP002076775<br/>           see abstract<br/>           &amp; JP 02 117 915 A (MITSUBISHI<br/>           PETROCHEMICAL CO LTD) 2 May 1990</p> | 1                     |
| X  | <p>DE 23 32 089 A (CIBA-GEIGY AG) 3 January<br/>           1974<br/>           see compounds 28, 29, 33, 37-39, 46</p>  | 1                     |
| X  | <p>CH 540 958 A (CIBA-GEIGY AG) 15 October<br/>           1973<br/>           see compounds 11, 13-26</p>   | 1                     |
| X  | <p>DE 23 64 396 A (BASF AG) 10 July 1975<br/>           see examples 1-3, 9-12, 16</p>  | 1                     |
| A  | <p>EP 0 287 909 A (SALUTAR INC) 26 October<br/>           1988<br/>           see page 3, lines 50-56; page 8, line 56 -<br/>           page 9, line 10; claims 1-10<br/>           &amp; US 5 039 511 A<br/>           cited in the application<br/>           &amp; US 4 933 156 A<br/>           cited in the application</p>  | 1,2,5-11              |
| A  | <p>V.K. SOOD ET AL: APPL. RADIAT. ISOT.,<br/>           vol. 48, no. 1, 1997,<br/>           pages 27-35, XP004015069<br/>           see page 28, experimental</p>  | 1,4                   |

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/07889

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 6-12 and 13-15  
because they relate to subject matter not required to be searched by this Authority, namely:  
see FURTHER INFORMATION sheet PCT/ISA/210
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

**FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210**

Although claims 6-12 are directed to a diagnostic method practised on the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

Although claims 13-15 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

In view of the vast number of theoretically conceivable compounds resulting from the combination of all claimed substituents the search had to be restricted for economic reasons (see Guidelines, Chapter III, paragraph 2.3). The search was limited to compounds of formula I, wherein Q represents structure IA.



# INTERNATIONAL SEARCH REPORT

Inter national Application No

PCT/US 98/07889

| Patent document<br>cited in search report | Publication<br>date | Patent family<br>member(s)   | Publication<br>date  |
|---|---------------------|--|--|
| WO 9634853 A                              | 07-11-1996          | AU 5669296 A<br>CA 2219880 A<br>EP 0823894 A<br>NO 975002 A  | 21-11-1996<br>07-11-1996<br>18-02-1998<br>02-01-1998   |
| CH 554821 A                               | 15-10-1974          | AT 299171 B<br>AT 288306 B<br>BE 721754 A<br>BE 748017 A<br>CH 554848 A<br>CH 513785 A<br>CS 174800 B<br>CS 174763 B<br>DE 1794386 A<br>DE 2012779 A<br>DE 1793482 A<br>DK 139845 B<br>FI 50238 B<br>FI 50238 C<br>FR 2044704 A<br>FR 1583595 A<br>GB 1247934 A<br>GB 1298577 A<br>HK 53076 A<br>NL 6814089 A,B<br>NL 7004439 A<br>SE 367399 B<br>US 3984399 A | 15-05-1972<br>15-01-1971<br>02-04-1969<br>28-09-1970<br>15-10-1974<br>15-10-1971<br>29-04-1977<br>29-04-1977<br>22-11-1973<br>25-02-1971<br>09-03-1972<br>30-04-1979<br>30-09-1975<br>12-01-1976<br>26-02-1971<br>14-11-1969<br>29-09-1971<br>06-12-1972<br>03-09-1976<br>09-04-1969<br>29-09-1970<br>27-05-1974<br>05-10-1976 |
| US 3755446 A                              | 28-08-1973          | BE 771239 A<br>CA 945553 A<br>DE 2039993 A<br>FR 2104174 A<br>GB 1351489 A<br>NL 7110945 A   | 14-02-1972<br>16-04-1974<br>17-02-1972<br>14-04-1972<br>01-05-1974<br>15-02-1972   |
| DE 2736889 A                              | 23-02-1978          | LU 75628 A<br>BE 857895 A<br>DK 367877 A<br>FR 2375191 A   | 13-04-1978<br>20-02-1978<br>20-02-1978<br>21-07-1978   |

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 98/07889

| Patent document<br>cited in search report | Publication<br>date | Patent family<br>member(s)   | Publication<br>date  |
|---|---------------------|--|--|
| DE 2736889 A                              |                     | JP 53025501 A<br>NL 7709102 A<br>SE 7709325 A  | 09-03-1978<br>21-02-1978<br>20-02-1978   |
| DE 2332089 A                              | 03-01-1974          | CH 574950 A<br>FR 2190815 A<br>GB 1391882 A<br>JP 1172503 C<br>JP 49058133 A<br>JP 58005213 B<br>US 4016156 A                | 30-04-1976<br>01-02-1974<br>23-04-1975<br>17-10-1983<br>05-06-1974<br>29-01-1983<br>05-04-1977               |
| CH 540958 A                               | 15-10-1973          | NONE   |  |
| DE 2364396 A                              | 10-07-1975          | CH 585692 A<br>DK 669774 A<br>FR 2255288 A<br>JP 50096622 A<br>US 4009193 A  | 15-03-1977<br>25-08-1975<br>18-07-1975<br>31-07-1975<br>22-02-1977   |
| EP 287909 A                               | 26-10-1988          | AU 620327 B<br>AU 1441988 A<br>CA 1302403 A<br>DE 3874187 A<br>JP 63290858 A<br>US 4933156 A<br>US 5008099 A<br>US 5039511 A | 20-02-1992<br>13-10-1988<br>02-06-1992<br>08-10-1992<br>28-11-1988<br>12-06-1990<br>16-04-1991<br>13-08-1991 |